



Figs. 23. Geographical distributions of *C. yamaokai* (●) and *C. nawai* (■) in Japan.

darker anterior portion of head. However, comparing nine syntypes of var. *atrigenatus* and six syntypes of var. *inconstans* with the nominate species, we have concluded that those are not significantly different from the latter in morphology.

Acknowledgments

We would like to thank Messrs. S. KUBOTA (Tokyo), A. KAWAZOE (Kuwana City), F. ITOH (Hokkaido Univ.) and K. MURATA (Utsunomiya Univ.), for their kindness in offering valuable materials, and to Dr. M. BRANCUCCI (Naturhistorisches Museum, Basel) for the loan of type materials. We also acknowledge our indebtedness to Dr. K. HAGA (Univ. of Tsukuba) for his helping the improvements of the manuscript and offering useful comments.

References

- EMERY, C., 1925. Hymenoptera, Formicidae, Formicinae. In WYTSMAN, M. P. (ed.), *Genera Insectorum*, fasc. 183: 1–302. La Haye.
- FOREL, A., 1912. Quelques fourmis de Tokio. *Annls. Soc. ent. Belg.*, 56: 339–342.
- ITO, T., 1912. *Camponotus Itoi* (*Myrmamblys*) FOREL stirps *tokioensis* ITO nov. st. In FOREL, A., *Quelques fourmis de Tokio*. *Annls. Soc. ent. Belg.*, 56: 341–342.
- 1914. Formicidarum japonicarum species novae vel minus cognitae. *Ibid.*, 58: 40–45.
- MENOZZI, C., 1940. Contribution a la fauna myrmecologique du Japon. *Mushi, Fukuoka*, 13: 11–