

*Beach Co.*: Boynton Beach (Wood & Davidson); *Sarasota Co.*: 30mi SE Sarasota, T38S, R22E, S.33 (J. T. Longino); Long Branch Key, Sarasota (A. C. Cole); Sarasota (A. C. Cole); *Seminole Co.*: Sanford (A. B. Gahan).

*P. cubensis*, mainland

ARGENTINA *Misiones*: El Dorado (N. Kusnezov); Esperanza (N. Kusnezov); *Salta*: Coronel Moldes (Monros).

BOLIVIA *Beni*: 14 km SW San Borja, 210 m (P. S. Ward); Espíritu (F. Koya).

BRAZIL *AM*: Ig. Marianil, Rio Branco Rd., 24 km NE Manaus (W. L. Brown); Manaus to Itacoatiara Hwy, km. 34 (W. L. & D. E. Brown); *MT*: Vila Vera (M. Alvarenga); *PA*: C. Araguaia (J. A. Rafael); Tucuruí (W. L. Overal); *PE*: Ilha Fernando Noronha (M. Alvarenga); *RJ*: Monumento Rodoviário (c.u.).

COLOMBIA *Cauca*: Popayán (W. & E. MacKay); Medellin, 1800 m (N. A. Weber); *Magdalena*: Gairaca Bay, Parque Tayrona, 0–5 m (C. Kugler); *Meta*: Pto. López (W. & E. MacKay).

COSTA RICA "Costa Rica" (F. Nevermann); *Alajuela*: 11 mi N Florencia (D. H. Janzen); *Guanacaste*: Est. Biol. Palo Verde, 10 m (J. T. Longino); N side Lago de Arenal, 570 m (J. T. Longino); Palo Verde (C. M. Herrera); *Heredia*: 3km S Pto. Viejo, 50 m (J. T. Longino); *Limon*: 5 mi. W Guapiles (D. H. Janzen); Hamburg Farm, Santa Clara (F. Nevermann); *Puntarenas*: 1 km NE Tárcoles, 20 m (P. S. Ward); Monteverde, 1100 m (P. S. Ward); Ojo de Agua, 800 m (J. T. Longino); Sirena, Corcovado Natl. Pk., 0–100 m (J. T. Longino); Sirena, Corcovado Natl. Pk., 50 m (J. T. Longino); *San José*: San Isidro de General (C. L. Hogue); San José (H. Schmidt).

GUATEMALA *Escuintla*: Escuintla (W. M. Wheeler); *Guatemala*: hills above Lake Amatitlán (D. H. Janzen); *Izabal*: Lago Izabal, 1.5 km NE El Estor (D. H. Janzen).

GUYANA *Mazaruni-Potaro*: Kartabo, Bartica Dist. (W. M. Wheeler); Kartabo Point (W. Steiner *et al.*).

MEXICO *Camp*: 29 mi E, 12 mi S Campeche (*Ruinas Edzna*) (D. H. Janzen); *Gro*: 25.4 mi S Chilpancingo (D. H. Janzen); *Is. Tres Marias*: Maria Madre I. (H. H. Keifer); Maria Magdalena I. (R. R. Snelling); *Jal*: Est. Biología Chamela, 100 m (P. S. Ward); *Mich*: 9 mi E Capirio, 800 ft. (E. Fisher & P. Sullivan); *Mor*: 5 mi S Cuernavaca (W. S. Ross); *Sin*: 20 mi E Villa Union, 235 m (E. I. Schlinger); *Tab*: 3 mi W Cárdenas (D. H. Janzen); 13 km W border Chiapas, Rt. 186 (W. P. MacKay); *Ver*: 9 km NNW Sontecomapan, 20 m (P. S. Ward); 10 km NW Isla, Ruta 145, 10 m (W. P. MacKay); Est. Biol. "Los Tuxtlas", nr. San Andres Tuxtla (G. Ibarra M.); Los Tuxtlas, 10 km NNW Sontecomapan, 200 m (P. S. Ward); Mirador (E. Skwarra); Palma Sola (P. Reyes); Sontecomapan (D. H. Janzen).

PANAMA *Canal Zone*: vic. Pta de los Chivos, 3 km SW Gatun, 100 m (W. L. Brown *et al.*).

PERU *Loreto*: Yurac, 67 mi E Tingo María (E. I. Schlinger & E. S. Ross).

VENEZUELA *Aragua*: El Limón, 480 m (C. J. Rosales); *Barinas*: Pueblo Nuevo (H. Romero); 17 km SSW Ciudad Bolívar, 240 m (P. S. Ward); *Cojedes*: 20 km W San Carlos (W. L. & D. E. Brown); *Monagas*: Caripe (EMC, JL, CHFH); Uverito (C. J. Rosales & J. A. Gonzales).

*Pseudomyrmex curacaensis* (Forel)

(Figures 10a, 10b, 26, 27)

*Pseudomyrma dolichopsis* var. *curacaensis* Forel 1912: 29. Two syntype workers, Curaçao (Bugnion) (MHNG) [Examined]. One syntype here designated LECTOTYPE.

*Pseudomyrmex curacaensis* [sic] (Forel); Kempf, 1961: 379 [Raised to species].

**Worker measurements (n=37).**— HL 0.85–1.21, HW 0.74–0.89, MFC 0.014–0.037, CI 0.70–0.88, OI 0.53–0.59, REL 0.51–0.58, REL2 0.64–0.77, OOI (-0.04)–0.47, VI 0.74–0.86, FCI 0.018–0.044, SI 0.44–0.49, SI2 0.60–0.74, FI 0.42–0.49, PDI 1.09–1.60, MPI 0.030–0.059, NI 0.56–0.69, PLI 0.71–0.89, PWI 0.59–0.78, PPWI 1.15–1.64.

**Worker diagnosis.**— Similar to *P. cretus* (*q.v.*); averaging smaller in size (HW 0.74–0.89), with longer head, shorter scapes, and longer eyes (Fig. 10a) (SI2 0.60–0.74; REL2 0.64–0.77) (see also Figs. 26 and 27); occipital margin varying from flat to (frequently) rather conspicuously concave, in full-face frontal view; funicular segments II and III about as long as broad, or nearly so (FLII 1.55–2.04, n=6). Propodeum and petiole (Fig. 10b) similar to that of *P. cretus*, except correspondingly more slender in smaller individuals. Sculpture, pilosity, and pubescence more or less as in *P. cretus*. Color variable: typically medium to dark brown, with lighter appendages, and a darker infuscated patch on the upper third of the