

- 439, 446
bequaerti (Enzmann),
Pseudomyrmex, 437
bequaerti (Wheeler),
Pseudomyrmex, 437, 446
bicinctus (Santschi),
Pseudomyrmex, 404
bicolor (Guérin), *Pseudomyrmex*,
404
biconvexus (Forel),
Pseudomyrmex, 438
biolleyi (Enzmann),
Pseudomyrmex, 445
boopis (Roger), *Pseudomyrmex*,
405
boxi (Wheeler), *Pseudomyrmex*,
444–445, 447
bradleyi (Enzmann),
Pseudomyrmex, 444
bradleyi (Wheeler),
Pseudomyrmex, 444, 447
browni Kempf, *Pseudomyrmex*,
404
brunneus (F. Smith),
Pseudomyrmex, 404
brunnipes (Enzmann),
Pseudomyrmex, 405
caeciliae (Forel), *Pseudomyrmex*,
404–406, 409–411, 418–420,
428, 448
canescens (F. Smith),
Pseudomyrmex, 404
canescens (Wasmann),
Pseudomyrmex, 437, 446
caroli (Forel), *Pseudomyrmex*, 447
castus (Wheeler), *Pseudomyrmex*,
404
catappa, Terminalia, 420
cephalicus (F. Smith),
Pseudomyrmex, 438
championi (Forel),
Pseudomyrmex, 404
chodati (Forel), *Pseudomyrmex*,
406, 427–428
cholericus (Wheeler),
Pseudomyrmex, 440, 446
cladoicus (F. Smith),
Pseudomyrmex, 404
cocae (Santschi), *Pseudomyrmex*,
404
colei (Enzmann), *Pseudomyrmex*,
404
concolor (F. Smith),
Pseudomyrmex, 405, 435–436,
441, 444–446
confusior (Forel), *Pseudomyrmex*,
406, 426
convarians (Forel),
Pseudomyrmex, 444, 452
cordiae (Forel), *Pseudomyrmex*,
404
cordobensis (Forel),
Pseudomyrmex, 405
coronatus (Wheeler),
Pseudomyrmex, 405, 436
costaricensis (Enzmann),
Pseudomyrmex, 405
cretus sp. nov., *Pseudomyrmex*,
404, 406, 408, 412–413,
416–417, 447
crucians (Wheeler),
Pseudomyrmex, 440, 447
cubaensis (Forel), *Pseudomyrmex*,
404, 406, 409–411, 413–415,
420, 428
culantrillo, *Zanthoxylum*, 420
culmicola (Forel), *Pseudomyrmex*,
429, 433
cumingiana, *Triplaris*, 420
curacaensis (Forel),
Pseudomyrmex, 404, 406, 408,
413–414, 416–417, 423, 447
damnosus (Wheeler),
Pseudomyrmex, 435