

- spiculus*, *sp. nov.*, *Pseudomyrmex*, 405, 429–434, 448
spinicola (Emery),
Pseudomyrmex, 404, 435, 438, 443–444, 447
squamiferus (Emery),
Pseudomyrmex, 404
stans, *Tecoma*, 420
stolli (Forel), *Pseudomyrmex*, 443, 447
subater (Wheeler & Mann),
Pseudomyrmex, 405
subtilissimus (Emery),
Pseudomyrmex, 403, 405, 429–430, 432–433, 435, 438
symbioticus (Forel),
Pseudomyrmex, 405, 445
tachigaliae (Forel),
Pseudomyrmex, 403, 405, 435–436, 441, 444–445, 447
tandem (Forel), *Pseudomyrmex*, 406, 419
tenuis (Fabricius), *Pseudomyrmex*, 403, 405, 444, 447
tenuissimus (Emery),
Pseudomyrmex, 405, 429–433
terminalis (F. Smith),
Pseudomyrmex, 404
termitarius (F. Smith),
Pseudomyrmex, 405, 444
testaceus (F. Smith),
Pseudomyrmex, 442, 447
thoracicus (Norton),
Pseudomyrmex, 404
tigrinus (Wheeler),
Pseudomyrmex, 445, 447
trigonus (Enzmann),
Pseudomyrmex, 445
trinervis, *Baccharis*, 420
triplaridis (Forel), *Pseudomyrmex*, 405, 435–436, 444–445, 447
triplarinus (Weddell),
Pseudomyrmex, 405
Pseudomyrmex, 405, 435–436, 444–445
tuberculatus (Enzmann),
Pseudomyrmex, 406, 422
ulei (Forel), *Pseudomyrmex*, 445, 447
unicolor (F. Smith),
Pseudomyrmex, 404
urbanus (F. Smith),
Pseudomyrmex, 404–406, 409, 411, 415, 420, 427–428
variabilis (F. Smith),
Pseudomyrmex, 439, 446
velifer (Stitz), *Pseudomyrmex*, 404
veneficus Wheeler,
Pseudomyrmex, 404, 439
venustus (F. Smith),
Pseudomyrmex, 404
vesanus (Wheeler),
Pseudomyrmex, 437, 446
viduus (F. Smith), *Pseudomyrmex*, 403, 405, 435–436, 444–447
villosum *sp. nov.*, *Pseudomyrmex*, 405, 429–430, 432, 434
vinneni (Forel), *Pseudomyrmex*, 404
virgo (Santschi), *Pseudomyrmex*, 404
vistanus (Enzmann),
Pseudomyrmex, 404
volatilis (F. Smith),
Pseudomyrmex, 404
vojtoskii (Enzmann),
Pseudomyrmex, 405
wasmanni (Wheeler),
Pseudomyrmex, 437, 446
weberi (Enzmann),
Pseudomyrmex, 405
wessonii (Enzmann),
Pseudomyrmex, 406, 422–423
wheeleri (Enzmann),
Pseudomyrmex, 405