

Comments: Notable features of *P. prioris* are its small size and long basal face of propodeum. It differs from the otherwise similar *P. succinus* by its possession of a less dorsoventrally compressed head (compare LCI) and shorter petiole ( $PLI > 0.55$ ).

***Pseudomyrmex succinus*, new species**

(Figures 9, 18)

Holotype worker: Dominican amber from Palo Alto region, purchased by P. S. Ward from S. Hendrickson (#PSW-DA17) (MCZC).

Paratype workers: Three workers in three separate pieces of Dominican amber, purchased for MCZC (no further data) (#PSW-DA1, #PSW-DA2, and #PSW-DA9); two workers in two pieces of Dominican amber from Santiago region (#PSW-DA23, #PSW-DA24) (from a series of six pieces of amber labeled H-10-114, UCBC).

Paratype queen: Dominican amber, purchased for MCZC (no further data) (#PSW-DA5).

Additional material: one worker in Dominican amber from Santiago region (#PSW-DA25) (from a series of six pieces of amber labeled H-10-114, UCBC).

Measurements, holotype: HW 0.51, HL 0.84, HD ca. 0.37, EL 0.47, SL 0.21, FL 0.47, FW ca. 0.22, LHT 0.43, DPL 0.49, BF 0.37, DF 0.18, PL 0.36, PH 0.19, DPW ca. 0.18, PPL 0.26, PPW ca. 0.29.

Measurements, paratype workers (n=5 unless otherwise noted): HW 0.48–ca. 0.55 (n=4), HL 0.81–0.89, HD 0.36–0.42, EL 0.45–0.50, SL 0.21–0.23, FL ca. 0.40–0.49, FW 0.21–0.24, LHT ca. 0.40–0.50, DPL 0.50–0.57, PL 0.33–0.37, PH ca. 0.16–0.20, DPW ca. 0.16–0.18 (n=3), PPL ca. 0.29–0.33 (n=4), PPW ca. 0.28–0.32 (n=3).

Measurements, paratype queen: HW ca. 0.47, HL 0.95, HD 0.40, EL 0.50, SL 0.23, FL 0.44, FW 0.24, LHT 0.45, PL ca. 0.39, DPW ca. 0.19, PPL 0.35, PPW 0.35.