

Labidus

- coecus* (Latreille, 1802), wqm
praedator s. str. (Fr. Smith, 1858), wqm

Neivamyrmex

- agilis* Borgmeier, 1953, w
andrei (Emery, 1901), m
angulimandibulatus Watkins, 1974, m
cloosae (Forel, 1912), m
cornutus Watkins, 1975, w
diabolus (Forel, 1912), m
fallax Borgmeier, 1953, w
fumosus (Forel, 1913), m
fuscipennis (Wheeler, 1908), m
graciellae (Mann, 1926), w
guerini (Shuckard, 1840), m
halidayi (Shuckard, 1840), m
harrisi (Haldeman, 1852), wqm
impudens (Mann, 1922), w
inflatus Borgmeier, 1958, m
klugi distans Borgmeier, 1953, m
leonardi (Wheeler, 1915), w
longiscapus Borgmeier, 1953, m
macropterus Borgmeier, 1953, m
manni (Wheeler, 1914), w
melanocephalus (Emery, 1895), w
melsheimeri (Haldeman, 1852), m
minor (Cresson, 1872), m
nigrescens (Cresson, 1872), wqm
opacithorax (Emery, 1894), wqm
pauxillus (Wheeler, 1903), wq
pilosus mandibularis (M. R. Smith, 1942), m
pilosus mexicanus (Fr. Smith, 1859), wqm
rugulosus Borgmeier, 1953, w
spinolai (Westwood, 1842), wqm
spoliator (Forel, 1899), m
sumichrasti (Norton, 1868), w
swainsoni (Shuckard, 1840), m
texanus Watkins, 1972, wqm
tristis (Forel, 1901), m

Nomamyrmex

- esenbecki mordax* (Santschi, 1928), wm
esenbecki wilsoni (Santschi, 1920), wm
hartigi (Westwood, 1842), wm