



FIG. 31. *Oligomyrmex (Aeromyrma) jeanneli* Santschi. Outline of head of soldier. After Santschi.

FIG. 32. *Hyldris africanus* (Karawaiew). Outline of worker. After Karawaiew.

FIGS. 33, 34. *Oligomyrmex (Aeromyrma) debilis* Santschi. 33. Outline of head of soldier. 34. Wing of male. Both after Santschi.

FIG. 35. *Hyldris africanus* (Karawaiew). Outline of head of worker. After Karawaiew.

FIG. 36. *Oligomyrmex (Aeromyrma) debilis* Santschi. Outline of thorax and pedicel of soldier. After Santschi.

***Cardiocondyla emeryi* subspecies *mahdii* Karawaiew**

Cardiocondyla emeryi subspecies *mahdii* KARAWAIEW, 1911, Rev. Russe Ent., vol. 11, p. 8, worker.

Khartoum, Anglo-Egyptian Sudan, is the type locality, and I took a worker of this or the typical form at Port Sudan in this country in 1939. The thorax length of the worker is 0.52 mm., as in the typical form.

***Cardiocondyla brevispinosa*, new species**

WORKER: Length, 2.1 mm.; of thorax including neck, 0.63 mm. Head in front view, excluding mandibles, one and one-fourth times longer than broad behind eyes; occipital margin sharply impressed medially, corners rounded; sides feebly convex, slightly broader behind eyes than in front; anterior clypeal margin produced over the base of the mandibles, feebly concave in the middle; eyes large, closer to the anterior clypeal margin than to the occiput; mandibles stout, evenly convex on their outer margins, with fine teeth, of which the second basal is