

CONSPECTUS SPECIERUM GENERIS *TRIP LARIDIS*.

trique- trum ; facibus :	subrotundis				<i>T. Bonplandiana</i> †.
		ovatis.	basi tubi adnatæ, achenium vix æquantes.		<i>T. Felipensis</i> †.
Perianthii laci- niæ interiores :	liberæ, achenio longiores.				<i>T. Noli-tangere</i> †.
		lobis 3 brevi- bus interjectis	tubo vix adnatæ.		<i>T. surinamensis</i> Ch. et Sch.
separatæ. Perianthii laci- niæ interiores :	tubo e basi ad mediam longitudinem adnatæ.				<i>T. peruviana</i> F. et M.
		inferne tereti-tubulosæ.			<i>T. Guayaquilensis</i> †.
trialatum. Perianthii lacinia exteriores :	tubo breviores aut æquales	} tubo sejunctæ aut vix adnatæ. Perianthium fructiferum	} 6 centim. longum	} 2 centim. vix longum.	<i>T. Pappigiana</i> †. <i>T. brasiliana</i> Ch. et Sch.
} tubo longiores.	} perianthii tubo longiores.	} perianthii tu- bo longiores.	} ovatis (subtus tomentosis)	} oblongis (subtus pubescentibus.	<i>T. Martiana</i> F. et M. <i>T. tomentosa</i> †. <i>T. Cumingiana</i> F. et M. <i>T. Lindeniana</i> †.

ACHENIUM :