

PSYCHE

VOL. XXXVII.

SEPTEMBER, 1930

No. 3

TWO NEW ANTS FROM JAVA

BY GEORGE C. WHEELER AND ESTHER W. WHEELER

University of North Dakota

*Phaulomyrma*¹ new genus

♂ Head flattened, slightly emarginate behind. Eyes protruding, large (diameter about one-third the length of the head), and situated near the anterior border. Clypeus indistinct or large and somewhat convex. Antennæ 13-segmented. Thorax compressed. Legs slender; two spurs on the middle and hind tibiæ; metathoracic legs, long, extending beyond the tip of the abdomen. Wings hairy; fore wings large, with one or three reduced veins; the margins fringed with hairs; hind wings much smaller. Hairs abundant especially on the dorsum. Integument smooth or with scattered hairy punctures. Genitalia large and projecting. Sagittæ wide at the base. Subgenital plate bifurcated.

Type: *Phaulomyrma javana* new species.

This genus is to be distinguished from *Leptanilla* by the presence of wing veins and the unusually large genitalia.

It is our opinion that *Leptanilla tanit* Santschi should also be included in this genus.

Phaulomyrma javana new species

♂ Length—1.2 mm.

Head large, about one-fifth of the entire length, slightly longer than broad, somewhat narrowed behind; the posterior border emarginate.

¹ *Phaulos*, trifling, paltry + myrmex, ant.