Emeryella schmitti Forel, Figs. 9-14.—9, head in anterior view, ×57; 10, 11, 13, 14, four body hairs, ×72; 12, left mandible in anterior view, ×72.

**Ectatomma (E.) tuberculatum (Olivier), Figs. 15-26.—15, right antenna and base in anterior view, ×210; 16, head in anterior view, ×55; 17, body hair, ×107; 18, larva in side view, ×8; 19, head in posterior view showing attached planidium, ×55; 20, labrum in posterior view (right half only), ×95; 21, spinules of integument arranged in irregular pattern, ×185; 22, spinules of integument arranged in short rows, ×185; 23, labrum and hypopharynx in anterior view (left half only), ×95; 24, left mandible in anterior view, ×82; 25, left maxilla in anterior view, ×95; 26, midventral surface of meso- and metathorax showing leg vestiges and alveoli of hairs, ×44.

PLATE V. Ectatomma (Poneracantha) bispinosum Emery, Figs. 1-5.—1, head in anterior view, ×55; 2, right antenna in anterior view, ×235; 3, right mandible in anterior view, ×95; 4-5, two body hairs, ×95.

Ectatomma (Parectatomma) sp., Figs. 13-16.—13, right antenna in anterior view, ×235; 14, profile of larva, ×6; 15-16, two body hairs, ×95.

Ectatomma (Gnamptogenys) tortuolosum (F. Smith), Figs. 6-12.—6-7 and 9-12, six body hairs, ×95; 8, profile of larva, ×6.

Proceratium crassicorne Emery, Figs. 17-19.—17, abdominal segment II in dorsal view, ×23; 18, abdominal segment IV to end of body in ventral view, ×23; 19, larva in side view, ×23.

Proceratium croceum (Roger), Figs. 20-27.—20, head in anterior view, ×59; 21, a dorsal boss in side view, ×167; 22, right mandible in anterior view, ×89; 3, body hair, ×370; 24, very young larva in ventral view, ×23; 25, very young larva in side view, ×23; 26, half grown larva in side view, ×23; 27, mature larva in side view, ×23.