



FIG. 1.—*Stigmatomma pallipes*: a, left mandible, anterior view, $\times 185$; b, larva, side view, $\times 12$. FIG. 2.—*Myopopone castanea*: a, head, anterior view, $\times 47$; b, left mandible, anterior view, $\times 111$; c, larva, side view, $\times 6$; d, labrum (right half, posterior view; left half, anterior view), $\times 139$; e, body hair, $\times 139$. FIG. 3.—*Paraponera clavata*: larva, side view, $\times 4$. FIG. 4.—*Prionopelta opaca*: immature larva, side view, $\times 22$. FIG. 5.—*Paranomopone relicta*: a, head in anterior view, $\times 67$; b and c, 2 types of body hairs, $\times 278$; d, larva, side view, $\times 15$; e, left labial palp, anterior view, $\times 333$; f, left mandible, anterior view, $\times 139$; g, left maxillary palp, anterior view, $\times 333$. FIG. 6.—*Typhlomyrmex pusillus*: a, right mandible, anterior view, $\times 235$; b, larva, side view, $\times 67$. FIG. 7.—*Acanthoponera (Anacanthoponera)* sp.: a, head, anterior view, $\times 71$; b-d, 3 types of body hairs, $\times 278$; e, left mandible, anterior view, $\times 139$. FIG. 8.—*Rhytidoponera tasmaniensis*: a, egg, $\times 16$; b, first-instar larva, side view, $\times 16$; c, second-instar larva, side view, $\times 16$; d, young larva, side view, $\times 16$; e, entangled hairs on dorsa of 2 larvae, $\times 96$. FIG. 9.—*R. mayri*: a, sexual (?) larva, side view, $\times 8$; b-d, 3 types of body hairs, $\times 85$. FIG. 10.—*R. cristata*: a-c, 3 types of body hairs, $\times 44$. FIG. 11.—*R. cerastes*: a, hooked hair of very young larva, $\times 85$, and tip (enlarged), $\times 350$; b, body hair of mature larva with frayed tip, $\times 85$, and tip (enlarged) $\times 350$; c and d, simple and bifid hairs of mature larva, $\times 85$; e, very young larva, side view, $\times 28$.