

Genus SOLENOPSIS Westwood

Markin et al. (1972: 1053): Life cycle of *Solenopsis invicta* Buren in an incipient colony: egg 6-8 days, larva 14-15 days, pupa 20-24 days.

Genus TETRAMORIUM Mayr

Tetramorium caespitum (Linnaeus). Donisthorpe (1927: 197): "The larvae were fed with disgorged liquid food as long as they were young and gathered together in groups, but when they grew older and were separated, the workers fed them with solid substances." Many larvae were hung on to the plaster walls of the nest by their anchor-tipped hairs.

LITERATURE CITED

- BERNARD, F.
1951. Super-famille des Formicoidea. *Traité de Zoologie, Tome X, Fasc. II*: 907-1119, 1258-1263, 1272-1275.
- CLOUDSLEY-THOMPSON, J. L.
1962. A note on the association between *Bengalia* spp. (Dipt., Calliphoridae) and ants in the Sudan. *Entomol. Monthly Mag.* 98: 177-179.
- DELAGE, BERNADETTE.
1968a. Recherches sur la fourmis moissonneuses du Bassin Aquitain: écologie et biologie. *Bull. Biol.* 100: 315-367.
1968b. Recherches sur les fourmis moissonneuses du Bassin Aquitain: éthologie. *Physiologie de l'alimentation. Ann. Sci. Nat., Zool. Biol. Anim.* (12)10: 197-265.
1972a. Une fourmi de Côte-d'Ivoire: *Melissotarsus titubans* Del., n. sp. *Insectes Sociaux* 19: 213-226.
1972b. Le polymorphisme larvaire chez les fourmis *Nematocrema* d'Afrique. *Insectes Sociaux* 19: 257-277.
- DONISTHORPE, H. ST. J. K.
1927. *British ants*. Geo. Routledge & Sons, London. 436 pp.
- ETTERSHANK, G.
1966. A generic revision of the world Myrmicinae related to *Solenopsis* and *Pheidologeton*. *Australian J. Zool.* 14: 73-171.
- JANZEN, D. H.
1966. Coevolution of mutualism between ants and acacias in Central America. *Evolution* 20: 249-275.
1967. Interaction of the bull's-horn acacia (*Acacia cornigera* L.) with an ant inhabitant (*Pseudomyrmex ferruginea* F. Smith) in eastern Mexico. *Univ. Kansas Sci. Bull.* 47: 315-558.
- KEMPF, W. W.
1972. A study of some Neotropical ants of genus *Pheidole* Westwood. I. *Studia Entomol.* 15: 449-464.