

CAUTION: Hairs have been omitted from most drawings of larvae in side view. In *Aphomomyrmex* they become too fine to be shown accurately at this magnification. In *Paratrechina* 5d most hairs have been broken off. In *Dendromyrmex* 8a they are too numerous.

LITERATURE CITED

- Cagniant, H. 1980. Études des stades larvaires, de la lignée des ailes et de la lignée des ouvrières dans des colonies avec reine et des colonies sans reine chez la fourmi *Cataglyphis cursor* Fonsc. (H. F.) Bull. Soc. Hist. Nat. Toulouse 116:192-206.
- Hinton, H. E. 1951. Myrmecophilous Lycaenidae and other Lepidoptera—a summary. Proc. South London Ent. and Nat. Hist. Soc. 1949-1950, pp. 111-175.
- Hölldobler, B. and E. O. Wilson. 1983. The evolution of communal nest-weaving in ants. Amer. Sci. 71:490-499.
- Kôriba, O. 1963. A parasitic life of *Polyrhachis lamellidens* F. Smith. First report. Kontyû 31:200-206.
- Wheeler, G. C. and Jeanette Wheeler. 1953. The ant larvae of the subfamily Formicinae. Ann. Entomol. Soc. Amer. 46:126-171, 175-217.
- Wheeler, G. C. and Jeanette Wheeler. 1976. Ant larvae: review and synthesis. Mem. Ent. Soc. Washington No. 7, 108 pp.
- Wheeler, G. C. and Jeanette Wheeler. 1982. Supplementary studies on ant larvae: Formicinae. Psyche 89:175-181.

Received June 6, 1985; accepted August 19, 1985.