


mandibles and antennal funiculi darker. Thorax and epinotum clouded with brown. Each gastric segment is crossed near its posterior edge by a dark band, which is narrow in the middle and as broad as the segment on either side where it surrounds a deep yellow spot. Venter yellow in the middle. Wings whitish hyaline, with brownish yellow veins and stigma.

Described from one female and many workers taken May 18 and 23 on the keys along the northern shore of the Southern Bight, Andros Island. One of the nests was in a dead branch of a gum mastic tree, the others in Tillandsias on mangroves at the edge of the 'swashes.'

28. **Cremastogaster victima** F. Smith var. **steinheili** Forel.—Numerous colonies containing males and winged females were found nesting in hollow twigs, and especially in the Tillandsias growing on the mangroves and other shrubs in the 'swashes' of Andros. These colonies were particularly abundant on Big Wood Key (May 16), Pot Key, and other keys along the course of the Southern Bight (May 18 and 19). A few specimens were also taken at West Bay on New Providence Island. All the specimens collected represent a form assignable to the var. *steinheili*, although the females are considerably larger (6–7 mm.) than a female type (4.6 mm.) of *steinheili* from Kingston, Jamaica (Forel), in my collection.



Macromischa pastinifera Emery var. *opacipes* var. nov. Worker.

29. **Macromischa pastinifera** Emery var. **opacipes** var. nov.

Two workers (Fig. I), found running over the ground under some dead palmetto leaves on a key in Crawl Creek, western Andros, agree with Emery's description and figure of the type from the "Bahama Islands," except in having the legs entirely opaque and finely punctate, instead of shining. On this account the specimens may be regarded as representing a distinct variety.

30. **Macromischa lucayensis** Forel. — This species, described by Forel from a single winged female from the "Bahama Islands," is