

Burmese form from a locality between Sablon and Naguillan, Luzon (H. M. Smith).

17. **Solenopsis geminata** (Fabr.) var. **rufa** (Jerdon). Numerous workers from Bais (L. E. Griffin) are recognizable as belonging to this variety on account of their red color and the spine on the mesosternum. Emery cites *S. geminata* from Manila and Antipolo, and it is probable that his specimens also belonged to this South Asiatic variety and not to the typical form of the species, which lives in the West Indies and the adjacent American tropics.

18. **Pheidologeton diversus** (Jerdon). Recorded by Smith under the name *Pheidole ocellifera* from "Burmah, Hong Kong, Philippine Islands." It is also cited by Ashmead from the locality last mentioned.

19. **Pheidologeton pygmaeus** Emery var. **albipes** Emery. Originally described by Emery from worker specimens taken by Simon at Antipolo, Luzon.

20. **Pheidole simoni** Emery. Described by Emery from a single soldier taken at Manila. It resembles *Ph. sauteri* from Formosa (*vide supra*) very closely, but has longer antennal scapes and the base of the gaster is opaque.

21. **Pheidole** sp. A form "allied to *Ph. longicornis* and *velox* Emery," of which Emery saw only a single worker from Antipolo.

22. **Cremastogaster ochracea** Mayr. Recorded by Emery from Antipolo, Luzon.

23. **Cremastogaster simoni** Emery. Originally described from worker specimens taken by Simon at Manila and Antipolo.

24. **Cremastogaster semperi** Emery. Like the preceding originally described from specimens taken at Manila by Simon.

25. **Cremastogaster longiclava** Emery. The types of this species were taken by Simon at Antipolo.

26. **Cremastogaster crassicornis** Emery. Types from Manila, collected by Simon. These resemble *C. brevis* Emery of Java.

27. **Cremastogaster bicolor** Mayr var. **imbellis** Emery. Described from single worker and female specimens taken at Manila and Antipolo by Simon.

28. **Tetramorium pacificum** Mayr subsp. **subscabrum** Emery. The worker of this subspecies was originally described from Antipolo, but, according to Emery, it occurs also in Ceylon and New Caledonia.

29. **Tetramorium guineense** (Fabr.). Numerous workers and dealated females of this common tropicolitan form were taken at Bais by L. E. Griffin. It is also recorded from the Philippines by Ashmead.

30. **Myrmicaria subcarinata** (F. Smith). A single worker taken at Benguet, Luzon (H. M. Smith) belongs to this species but differs from