

Kingston (Grabham); workers and females from Mandeville and Balaclava (Wight).

16. **Cardiocondyla emeryi** Forel.—Several workers from Balaclava and Troy (Wight).

17. **Solenopsis geminata** (Fabr.).—Numerous specimens of all three phases of the typical dark form of the species from Troy and Balaclava (Wight), and Kingston (Grabham and Andrews).

18. ***Solenopsis azteca** Forel.—Several workers from Mandeville (Wight) agree closely with a type specimen of this species from St. Vincent in my collection.

19. **Solenopsis azteca** var. **pallida** Wheeler.—Numerous workers from Mandeville and Balaclava (Wight).

20. ***Solenopsis hammari** Mayr?—A few workers from Mandeville (Wight) agree very closely with Emery's figure of this species (Bull. Soc. Ent. Ital., XXXVII, 1905, fig. 11) and probably belong either to it or to a dark subspecies of *S. decipiens* Emery. The antennal scapes are rather short and the clypeal teeth stout as in *hammari*, but the shape of the epinotum in profile is more like that of *S. decipiens* subsp. *abjecta* Emery, *i. e.*, with a more pronounced mesoëpinotal constriction than in *hammari*. The specimens measure about 1.5 mm. and are dark brown, with the mandibles, legs and gaster, except the first segment, yellow.

21. **Pheidole fallax** Mayr.—Soldiers and workers from several colonies; Balaclava (Wight); Kingston (Wight and Grabham).

22. **Pheidole megacephala** (Fabr.)—Soldiers and workers from Mandeville (Wight) and Kingston (Grabham).

23. **Pheidole subarmata** Mayr.—Soldiers and workers from several colonies; Balaclava (Wight).

24. **Pheidole radoszkowskii** Mayr var. **opacissima** Forel.—A few workers from Kingston (Grabham) agreeing very closely with topotypes received from Prof. Forel.

25. ***Pheidole caribbæa** sp. nov.

Soldier. Length 3.5–4.3 mm.

Head rather small, without the mandibles a little longer than broad, as broad in front as behind, convex above and below, with rounded sides and posterior corners and rather deep, angular occipital excision, which is continued forward over the vertex as a well-defined median groove to the frontal area. Eyes moderately large, convex, in front of the middle of the sides of the head. Mandibles convex, with two stout apical and no basal teeth. Clypeus short, flattened and ecarinate in the middle, its anterior border with a median excision. Frontal area subtriangular, rounded behind, as long as broad, impressed, without a median carinula. Frontal