

11. *Tetramorium guineense* (Fabricius)

*Formica guineensis* Fabricius, Ene. Syst., 2, 1793, p. 357 ♀.

*Myrmica bicarinata* Nylander, Acta. Soc. Sc. Fennic. 2, 1846, p. 1061 ♀ ♀.

*Myrmica cariniceps* Guérin, Rev. Mag. Zool. (2) 4, 1852, p. 79 ♀.

*Myrmica kollari* Mayr. Verh. zool. bot. Ver. Wien, 5, 1855, p. 425 ♀ ♀ ♂.

*Tetramorium cariniceps* Roger, Berlin, Ent. Zeitschr. 5, 1861, p. 171.

*Myrmica reticulata* F. Smith, Trans. Ent. Soc. London (3) 1, 1862, p. 33 ♀.

*Tetramorium kollari* White, Ants and Their Ways, 1883, p. 262 ♀.

*Tetramorium guineense* Mayr, Verh. Zool. bot. Ges. Wien, 20, 1870, p. 972 ♀.

A single worker of this undoubtedly introduced tropicopolitan ant from South Albemarle (F. X. Williams) and 14 workers taken by the "Albatross" (U. S. Nat. Mus.) but bearing only the label "Galapagos Islands, 1899." Emery records the species as taken also on the boat at Chatham Island by Dr. G. Baur.

12. *Tetramorium (Tetrogmus) simillimum* (F. Smith)

*Myrmica simillima* F. Smith, List Brit. Anim. Brit. Mus. 6, 1851, p. 118 ♀.

*Tetrogmus caldarius* Roger, Berl. Ent. Zeitschr. 1, 1857, p. 12 ♀ ♀.

*Myrmica (Leptothorax) simillima* F. Smith, Cat. Brit. Fos- sor. Hymen., 1858, p. 31 ♀.

*Myrmica caldaria* Meinert, Naturv. Aft. Dansk. Vid. Selsk. (5) 5, 1860, p. 334 ♀ ♀ ♂.

*Tetramorium simillimum* Mayr, Europ. Formicid. 1861, p. 61 ♀.

A single worker taken by Dr. Williams on Charles Island, "under stones near top of crater." This species, like *T. guineense*, is a tropicopolitan "tramp" of Old World origin. It is not infrequently found in hot houses in temperate regions.