

- Odontomachus hæmatoda** L.—Port of Spain, Gasparee Island and Sangre Grande, (R. Thaxter), ♂, ♀; Botanical Garden, Port of Spain, (Wheeler), ♂.
- Odontomachus hæmatoda** subsp. **insularis** Guérin var. **hirsutiusculus** F. Smith.—Port of Spain, (R. Thaxter), ♂.
- Odontomachus hæmatoda** L. subsp. **meinerti** Forel.—Port of Spain, (R. Thaxter), ♂.

Pseudomyrmæ

- Pseudomyrma championi** Forel var. **paulina** Forel.—Port of Spain, (Aug. Busck; R. Thaxter), ♂.
- Pseudomyrma elegans** F. Smith subsp. **breviceps** Forel.—Savanna, (A. Forel), type-locality, ♂; Trinidad, (A. Busck), ♂; Botanical Garden, Port of Spain, (Wheeler), ♂.
- Pseudomyrma elongata** Mayr.—Trinidad, (A. Forel), ♂; Port of Spain, (P. B. Whelpley), ♂.
- Pseudomyrma flavidula** F. Smith.—Port of Spain, (P. B. Whelpley), ♂.
- Pseudomyrma gracilis** Fabr.—Port of Spain, (P. B. Whelpley), ♂; Botanical Garden, Port of Spain, (Wheeler), ♂.
- Pseudomyrma culmicola** Forel.—Trinidad, (A. Forel), type-locality, ♂.
- Pseudomyrma excavata** Mayr.—Port of Spain, (R. Thaxter), ♂; Botanical Garden, Port of Spain, (Wheeler), ♂.
- Pseudomyrma filiformis** Fabr.—Trinidad, (Urich), ♂.
- Pseudomyrma kuenckeli** Emery.—Trinidad, (Urich); Botanical Garden, Port of Spain, (Wheeler), ♀.
- Pseudomyrma nigropilosa** Emery subsp. **laticeps** Forel.—Trinidad, (A. Forel), type-locality, ♂.
- Pseudomyrma pallida** F. Smith.—Aripa Savanna, (R. Thaxter), ♂; Trinidad, (Urich), ♂.

Pseudomyrma icterica, new species

WORKER.—Length, nearly 6 mm.

Head subrectangular, about one-fifth longer than broad, as broad in front as behind, with feebly convex sides and feebly and broadly excised posterior border; in profile convex above and below. Mandibles moderately convex, with two larger apical and several minute basal teeth. Clypeus bluntly carinate behind, its border broadly sinuate on each side, with a distinct rectangular lobe, rounded at the corners. Frontal carinæ small, closely approximated; frontal groove feeble. Eyes moderately large and convex, about two-fifths the length of the sides of the head. Antennal scapes reaching the middle of the head; first funicular joint a little longer than broad, second broader than long, remaining joints slightly longer than broad, terminal joint longer. Thorax rather narrow; the pro- and mesonotum together a little longer than the epinotum; the mesoëpinotal constriction abrupt, rather long and deep, its im-