

- Camponotus (Myrmobrachys) auricomus** Roger.—Botanical Garden, Port of Spain, (Wheeler), ♀.
- Camponotus (Myrmobrachys) beebei** Wheeler.—Near Port of Spain, (R. Thaxter), ♀.
- Camponotus (Myrmobrachys) brettessi** Forel.—Trinidad, (A. Forel), ♀.
- Camponotus (Myrmobrachys) canescens** Mayr.—Botanical Garden, Port of Spain, (Wheeler), ♀.
- Camponotus (Myrmobrachys) excisus** Mayr.—Port of Spain, (R. Thaxter), ♀.

**Camponotus (Myrmobrachys) godmani** Forel variety **palliolatus**, new variety

Six minor workers taken near Port of Spain by Prof. R. Thaxter agree closely with Forel's description of the types from Mexico and Central America, except that the tibiæ, upper surfaces of the femora and upper surface of the thorax are dark brown; the pigment on the pronotum being aggregated in two large, rather indefinite spots. The posterior portion of the head seems also to be more extensively infuscated than in the typical form.

- Camponotus (Myrmobrachys) lindigi** Mayr.—Gasparee Island, (R. Thaxter), ♀.
- Camponotus (Myrmobrachys) senex** F. Smith.—Port of Spain, (R. Thaxter, P. B. Whelpley), ♀.
- Camponotus (Myrmobrachys) zoc** Forel.—Trinidad, (A. Forel); Ariopita Valley, (B. D. Chipman), ♀.
- Camponotus (Myrmamblys) novogrenadensis** Mayr.—Botanical Garden, Port of Spain, (Wheeler), ♀.
- Camponotus (Myrmocladæcus) rectangularis** Emery var. **setipes** Forel, —Trinidad, (Urich), type-locality, ♀.
- Camponotus (Myrmocladæcus) latangulus** Roger.—Port of Spain, (R. Thaxter), ♀.
- Camponotus (Myrmocladæcus) bidens** Mayr.—Port of Spain, (R. Thaxter), ♀.
- Camponotus (Pseudocolobopsis) claviscapus** Forel.—Trinidad, (Urich), type-locality, ♀, ♂.