

33. *Euponera* (*Mesoponera*) *marginata* ROGER. ♀ Churubamba and Mojos, Bolivia (N. HOLMGREN); Brazil (F. SAHLBERG). May.

The striæ on the meso- and metapleuræ are less pronounced than in Brazilian specimens but the differences do not seem to be sufficient to justify a varietal name.

34. *Euponera* (*Mesoponera*) *fawceti* EMEY. ♀♂♂ Lagunillas, Bolivia (N. HOLMGREN); Bolivia and Peru (STAUDINGER); Huadquina, Peru, 5000 ft. (Yale Peruvian Expedition); Santa Marta, Colombia (F. M. GAIGE).

Female (undescribed). Length 9—10.5 mm.

Very similar to the worker, except for the usual differences and the shape of the petiole, which is higher, broader and more compressed anteroposteriorly. The pilosity is longer and a little more abundant. The wings are distinctly infuscated, with brown veins and darker pterostigma.

Male (undescribed). Length 10 mm.

Slender. Head through the eyes much broader than long, the latter and the ocelli very large and convex. Cheeks very short. Mandibles very small, bluntly pointed, edentate. Clypeus with a large, blunt, projecting tubercle in the middle, so that its outline in profile is almost rectangular, its anterior border straight and transverse. Antennæ very long (nearly 8 mm.) and filiform; the scapes only about $\frac{1}{3}$ as long as the second funicular joint; first funicular joint not broader than long and not broader than the remaining joints. Mesonotum with the Mayrian furrows indistinct posteriorly. Epinotum sloping, its base somewhat shorter than its declivity. Petiole somewhat like that of the worker but lower, thicker and somewhat more conical. Gaster and legs slender.

Rather shining, especially the abdomen. Head and thorax indistinctly and unevenly punctate-rugulose; epinotal declivity more distinctly transversely rugulose, especially below. Gaster very finely and superficially shagreened.

Pilosity and pubescence less developed than in the worker, the hairs on the head and thorax darker and more flexuous.

Color as in the worker; genitalia and mouthparts, except the mandibles, brownish yellow. Wings as in the female.

Lives together with a podurid. Collected Sept. 12th.

35. *Euponera* (*Mesoponera*) *arhuaca* FOREL. ♀ San Firmin, Bolivia (N. HOLMGREN). Under a decaying stub.

36. *Euponera* (*Trachymesopus*) *stigma* FABR. ♀ Peru (N. HOLMGREN).

37. *Euponera* (*Trachymesopus*) *holmgreni* sp. nov.