

87. *Monomorium pharaonis* LINN. ♀♀. Lower Amazon; Fontebôa, Brazil (HAHNEL); São Paulo, Brazil (A. HEINE); São Leopoldo, (J. W. STAHL).
88. *Megalomyrmex balzani* EMERY. ♀ San Fermin, Bolivia (N. HOLMGREN); ♀♂ Songo and Mapiri, Bolivia; Pozuzu, Peru (STAUDINGER).

Male (undescribed). Length 8 mm.

Head through the eyes as broad as long, rounded behind, without distinct posterior corners, the occipital border raised, or marginate. Eyes large, cheeks very short. Mandibles well-developed, with oblique, coarsely 4-toothed apical borders. Clypeus convex and bluntly subcarinate in the middle. Antennæ very long and filiform; scapes nearly four times as long as broad and distinctly longer than the second funicular joint; first funicular joint as long as broad, succeeding joints long and cylindrical, increasing in length and decreasing in thickness towards the tip. Thorax large, through the eyes; mesonotum very convex and evenly rounded, hemispherical, without Mayrian furrows. Epinotum long and sloping, feebly convex in profile, without distinct base and declivity and without a median dorsal groove. Petiole about twice as long as high, the node almost obsolete; the node of the postpetiole also very low, longer than high and longer than broad, inverted heart-shaped when seen from above, the posterior border somewhat impressed in the middle. Gaster elongate-elliptical, convex above. External genital valves roundly triangular; cerci apparently absent. Legs rather long and slender. Wings with sharply defined veins, with a large discoidal, a single cubital and an open radial cell and with both branches of the cubitus well-developed.

Sculpture and color much as in the worker, but the pilosity is very feebly developed, almost lacking on the body, very short and appressed on the appendages. Wings hairy, distinctly infuscated, with dark brown veins and pterostigma.

89. *Megalomyrmex goeldii* FOREL. ♀♂ Nogueira, Brazil (T. BORGMEIER).

Male (undescribed). Length nearly 4 mm.

Head including the eyes, a little longer than broad, broad and rounded behind, with straight, marginate occipital border. Eyes rather small. Cheeks well-developed. Mandibles 5-toothed. Clypeus convex in the middle. Antennal scapes long, as long as joints 2—4 of the funiculus, the first joint about $1\frac{1}{2}$ times as long as broad; joints 2—12 cylindrical, increasing in length towards the tip. Thorax shaped much as in *balzani*,