

MYRMICINÆ.

Aphænogaster (*Deromyrma*) **feæ** Emery.

Female (undescribed). Length 6.5-7 mm.

Resembling the worker. Cheeks with two parallel carinæ, instead of one, forming the external border of the antennal fovea. Mesonotum not overarching the pronotum, convex, as broad as long. Epinotum sloping, the base twice as long as the declivity, feebly concave, the spines stout, fully twice as long as their basal diameter, as long as the declivity, directed upward and backward. Petiole and postpetiole as in the worker, but their nodes higher and stouter. Gaster more voluminous than the thorax.

Sculpture, pilosity and color as in the worker, but the base of the epinotum coarsely and regularly, transversely rugose, the petiolar node transversely rugulose. Wings long, their membranes uniformly brownish, their veins and pterostigma darker brown.

Male (undescribed). Length 4.5 mm.

Head resembling that of the worker but shorter, the portion behind the prominent eyes less convex and narrowing more rapidly to the collar. Mandibles with a sharp apical tooth and five or six subequal basal denticles. Cheeks short and straight, without carinæ. Clypeus convex, its anterior border rounded and entire. Antennal scapes reaching a little beyond the posterior ocelli; funiculi slender, first joint longer than broad and distinctly thicker than the succeeding joints. Mesonotum strongly overarching the very short pronotum, very convex in front; epinotum long, the base long and sloping, strongly concave in front, horizontal and feebly convex behind, without spines or teeth, but with a distinct angle on each side where it passes into the very short declivity. Petiole and postpetiole as in the worker, but the nodes lower and that of the petiole more angular in profile.

Sculpture and color as in the worker and female, very smooth and shining, but the base of the epinotum and upper surface of petiolar node without rugæ. Hairs sparser, more