

Hairs on the scapes and legs slightly longer and less appressed than in *lineolata*.

Head, thorax, pedicel, coxæ and femora and more or less of the venter of the first gastric segment bright yellowish red, more vivid than in the female of the var. *clara* Mayr of the subsp. *laeviuscula*. Gaster scutellum, metanotum and three large elongate spots on the metanotum, one anteromedian, the others parapsidal, jet black; mandibles, antennæ, tibiæ, knees and tarsi darker and more brownish red; bases and sometimes the whole of the scapes, the tips of the funicular clubs, the ventral portions of the mesosterna, the inner border of each ocellus and the posterior corners of the head, more or less blackened or infuscated. Wings whitish hyaline, the pterostigma and anterior veins of the forewings very faintly yellowish, the remaining veins colorless.

*Male*.—Length 4 mm.

Very similar to the male of *lineolata* but somewhat more slender; gaster, especially, much narrower and more elongate. Mandibles more linear and narrower, furnished with three more distinctly equal denticles. Funicular joints apparently a little more cylindrical and less monillate. Pronotum distinctly less convex anteriorly and less over-arching the pronotum. Epinotum more rounded with less distinct base and declivity and no indications of teeth, except in one of the six specimens.

Somewhat smoother and more shining than the male of *lineolata* and more like that of *laeviuscula*. Mesonotum very finely and rather indistinctly striated and sparsely punctate.

Hairs whitish, almost lacking on the head, thorax and gaster, where they are present, though sparse, in *lineolata*; gaster and legs merely with short, sparse, appressed pubescence; antennæ and genital appendages with short, erect or suberect, very dense white pubescence.

Jet black; mandibles, gaster, antennæ, legs and alar insertions dark brown. Wings whitish hyaline with the pterostigma and anterior veins of forewings even more feebly