

A rather rare ant, distributed throughout the United States and represented by several varieties or subspecies, some of which are still undescribed.

- (4) *Euponera (Trachymesopus) stigma* Fabr.—Lake Worth (J. Schmitt).

A common neotropical form, but known only from this single locality in the United States.

- (5) *Ponera coarctata* Latreille subsp. *pennsylvanica* Buckley—Monticello; Gainesville; cited from "Florida" by Emery.

Common and widely distributed in the states east of the Mississippi.

- (6) *Ponera opaciceps* Mayr—Royal Palm Park and Dunedin (W. S. Blatchley).

- (7) *Ponera trigona* Mayr var. *opacior* Forel—Jacksonville (Wheeler).

Like the preceding a widely distributed neotropical form.

- (8) *Leptogenys (Lobopelta) elongata* Buckley subsp. *manni* Wheeler—Bellair.

Apparently a local race of the typical *elongata*, which ranges from Georgia to Texas.

- (9) *Odontomachus hamatoda* L. subsp. *insularis* Guérin—Lake Worth (J. Schmitt); Enterprise (W. Beutenmueller); Biscayne Bay (Mrs. A. T. Slosson); Coconut Grove, Lower Matabonie Key and Paradise Key (Wheeler); Tallahassee; Gainesville; Monticello; Lakeland. Deep Lake; Miami (A. E. Wight); St. Petersburg (H. Roster); Royal Palm Park (W. S. Blatchley).

Very generally distributed throughout the state and living in small colonies under stones, logs, boards, etc. It seems to be an immigrant from the West Indies, where it is common.

Subfamily Pseudomyrmecinae

- (10) *Pseudomyrma Brunnea* F. Smith—Haw Creek, Volusia Co. (T. Pergande); Tallahassee; Royal Palm Park and Dunedin (W. S. Blatchley); Lakeland; Miami (H. Hebard); Paradise Key, Long Pine Key, Lower Matabonie Key and Pine Crest (Wheeler).