

- (27) *Pheidole metallescens* Emery—St. George, type locality (T. Pergande); Jacksonville (Wheeler); Royal Palm Park (W. S. Blatchley).
- (28) *Pheidole morrissi* Forel—Royal Palm Park (W. S. Blatchley); St. Petersburg (H. Raster); Inverness (C. M. Weed).
- (29) *Cardiocondyla emeryi* Forel—Miami (Amer. Mus. Nat. Hist.).
- (30) *Cardiocondyla nuda* Mayr var. *minutior* Forel—Miami (W. E. Wight), numerous specimens.

This tiny ant has not been recorded heretofore from the United States.

- (31) *Cardiocondyla wroughtoni* Forel var. *bimaculata* Wheeler.

Two colonies containing a number of females, taken in hollow culms of sedges at Royal Palm Park (Wheeler). This variety was originally described from Formosa and closely resembles the var. *hawaiiensis* Forel, except that the two spots on the sides of the first gastric segment of the worker are large and dark brown. Sometimes there is a third smaller and paler brown spot in the middorsal like. The typical *wroughtoni* was originally described from India. This ant, like the preceding, is new to the fauna of the United States and has probably been very recently introduced from the Orient in living plants.

- (32) *Xenomyrmex stollii* Forel subsp. *floridanus* Emery—Lake Worth (T. Pergande), nesting in a twig of a gumbo-limbo tree (*Sideroxylon masticodendron*); Dunedin, workers and females (W. S. Blatchley).
- (33) *Xenomyrmex stollii* subsp. *rufescens* Wheeler.

Recently described from a single dealated female taken on Long Pine Key (Wheeler).

- (34) *Crematogaster (Acrocoelia) ashmeadi* Mayr—Key West (T. Pergande); Miami (Wheeler, A. E. Wight, P. Laurent); Card's Point, Key Largo, Long Pine Key, Lower Matabombie Key, Pine Crest and Bottle Point Key, on red mangrove (Wheeler); Biscayne Bay (Mrs. A. T. Slosson), on mangrove trunks; St. Augustine (C. T. Brues), in flower-stalks of *Yucca*; Dunedin (W. S. Blatchley).