

Grove, Paradise Key, Long Pine Key, Planter on Key Largo (Wheeler); Lake Worth (J. Schmitt); Marco; Everglade; Royal Palm Park, Paradise Key (W. S. Blatchley); Gulfport (Reynolds).

These localities are all in the southern third of the peninsula. *C. tortuganus* forms rather small colonies under stones. In the "Genera Insectorum" Emery doubtfully cites it as a subspecies of *C. conspicuus* Smith, but it seems to me preferable to regard it as an independent species.

(72) *Camponotus (Tanæmyrmex) incensus* sp. nov.

WORKER MAJOR. Length about 7 mm.

Resembling *tortuganus* but much smaller; (the latter measures 10–11 mm.) head of the same shape but proportionally shorter, with more acute antero-lateral clypeal angles, more nearly straight, crenated anterior clypeal border, less impressed frontal area, posteriorly more approximated frontal carinæ and somewhat shorter and more compressed antennal scapes. Thorax distinctly shorter with more convex and more even dorsal curvature, broader through the pronotum, which is less flattened above; base of epinotum forming a more distinct angle with the declivity. Grooves of the hind tibiæ less distinct. Surface of body less sharply shagreened, so that the surface, especially of the occiput, posterior corners of the head and the thorax distinctly smoother and more shining. Cheeks and sides of head with sparse but shallow, elongate foveolæ or coarse punctures, occiput with a radiating series of linear punctures and dorsal surface of pronotum with several similar impressions. Pilosity as in *tortuganus*, but paler yellow and pubescent on legs shorter. Color similar but the thorax is paler and more yellowish and the occiput and posterior corners of the head are paler, reddish brown. Gaster brown, paler than in *tortuganus*, the segments with a narrow dark brown band near the pale posterior border.

WORKER MINOR. Length .63 mm.

Very similar to the minor of *tortuganus* but much smaller and paler, the epinotum distinctly more angulate in profile, the base straight and nearly concave, the declivity very short. Thorax paler than in *tortuganus*, front, clypeus and cheeks anteriorly yellow, mandibles brownish yellow; gaster colored like that of the major.

Described from a single major and two minor workers which I collected many years ago (May 6, 1904) on Pigeon Key, near Miami, Florida

- /c (73) *Camponotus (Tanaemyrmex) Sosius* Roger—Sanford (J. Schmitt); Green Cove Spring (Mrs. Mary Treat); Dundenin (W. S. Blatchley); De Funiak Springs; Estera; Tampa; Clearwater; Kissimee.