

This tropical ant, which is common in Cuba and Mexico, nests either in the hollow branches of trees, especially *Coccoloba uvifera*, in the stems of weeds or in Tillandsias. It is confined to the southern third of the peninsula.

- (80) *Camponotus (Colobopsis) impressus* Roger—Lake Worth J. Schmitt); Bellair (Mrs. A. T. Slosson); Dunedin and Royal Palm Park (W. S. Blatchley); Paradise Key, Long Pine Key and Pine Crest (Wheeler).

I found this ant very common in the culms of sedges. It nests precisely like *C. culmicola* Wheeler of the Bahamas. Males and winged females were present in most of the colonies during late December and early January.

- (81) *Camponotus (Colobopsis) pylartes* Wheeler.

Dr. M. R. Smith records this species as having been taken by W. S. Blatchley in Royal Palm Park.

- (82) *Paratrechina longicornis* Latr.—Miami (A. E. Wight); Coconut Grove (Wheeler); Royal Palm Park (W. S. Blatchley); St. Petersburg (H. Raster).

A common tropicopolitan and introduced species, the "crazy ant" ("hormiga loca") of the West Indies.

- (83) *Paratrechina (Nylanderia) parvula* Mayr—Dunedin (W. S. Blatchley). Recorded by Mayr and Emery from "Florida."

- (84) *Paratrechina (Nylanderia) vividula* Nylander—Gainesville; Miami (A. E. Wight).

- (85) *Paratrechina (Nylanderia) bourbonica* Forel var.—Miami Beach (S. O. Hill); Miami (A. E. Wight) Apr., May, 1924; Coconut Grove (Wheeler).

Of recent introduction from the Orient.

- 1 c (86) *Prenolepis imparis* Say var. *testamea* Emery—St. Augustine (C. T. Brues); Tallahassee.

Known only from the northern portion of the state; common as far north as New Jersey.

- (87) *Lasius niger* L. subsp. *alienus* Förster var. *americanus* Emery—Monticello. Mayr had already recorded this very common northern ant from "Florida."