

Nylanderia fulva nesiotis Wheeler

Four workers from Hood Island (IV.20.'32), one from Iguana Cove, Albemarle Island (V.21.'32); two winged females from Tagus Cove, Albemarle (V.25.'32; V.27.'32) and one from James Island (VI.4.'32). Previously known from James and Indefatigable islands.

REVILLAGIGEDO ISLANDS

Solenopsis geminata Fabr.

A single worker of the typical black form from Socorro Island (III.26.'32) and a very small and more reddish specimen from Clarion Island (III.24.'32).

Camponotus (Tanæmyrmex) picipes Oliv. var. ?

Two minima workers from Socorro Island, 2000 ft. (III.7.'32). They probably represent an undescribed variety of the Mexican *picipes*, but further determination is impossible without the major worker.

COCOS ISLAND

Camponotus (Myrmaphænus) cocosensis Wheeler

Seven workers, four females and six males (VI.28.'32).

The *female* (undescribed) measures 11–12 mm. and resembles the worker major, but the head is less narrowed anteriorly and the antennal scapes are longer. Thorax elongate-elliptical, slightly broader than the head; mesonotum as long as broad; base of epinotum convex, decidedly shorter than the subperpendicular, concave declivity. Superior border of petiolar node rather deeply and semicircularly excised. Surface of head and thorax much smoother and more shining than in the worker. Erect hairs on the head and thorax shorter and sparser. Head and appendages reddish yellow; thorax and petiole deep red. Wings long (12 mm.), yellow, with somewhat darker resin yellow veins and pterostigma.

The *male* (undescribed) measures 6.5–7 mm. Head through the eyes very nearly as wide as long; cheeks straight; clypeus bluntly carinate; mandibles triangular, with well-developed but edentate apical border; eyes and ocelli large and prominent; antennæ long and slender. Thorax stout, with large mesonotum, much broader than the head; epinotum short and convex, with subequal base and declivity. Petiolar node low and thick, its obtuse superior border broadly impressed in the middle. Legs long and slender. Pilosity pale, sparse and of uneven length as in the female. Head and thorax subopaque as in the worker; scutellum and epinotum smoother and more shining. Head, genitalia, appendages and sutures of thorax brownish or reddish yellow; remainder of thorax dark brown; gaster black. Wings yellow as in the female but both membranes and veins distinctly paler.