

nearly straight anteriorly, not carinate; disc of frontal carinae small, ciliate, broadly sulcate in the middle; frontal groove not reaching to the middle of the head. Eyes very minute, not distinctly composite. Antennal scape barely reaching the occipital border; funiculus gradually incrassated towards its tip; club indistinctly 5-jointed, all the joints except the first and last broader than long. Thorax stout, dorsum feebly arcuate, not impressed, basal and declivous surfaces of epinotum meeting at a very rounded angle; promesonotal suture distinct; mesoëpinotal suture feebly indicated or obsolete; mesonotum transverse, half as long as the pronotum, both, including the pleurae and all the sterna, punctate, slightly less opaque than the head above; epinotum shining, with separated piligerous punctures, slightly narrower than the pronotum, with concave sides, the declivous surface broad, transversely excavated for the reception of the petiolar scale, its sides marginate. Petiolar scale seen from above semilunar, or semicircular, convex anteriorly, nearly straight or transversely subconcave posteriorly, seen from the side nearly straight anteriorly or slightly concave, the posterior passing arcuately into the superior surface. Sculpture of the whole abdomen like that of the epinotum. Length 2.5–2.75 mm.

“Notable on account of the structure of the petiole and very minute eyes, in which separate facets are not with certainty discernible.

“Lemien (New Guinea), collected in siftings (L. Bíró).”

***Selenopone clavicornis* (Emery)**

EMERY, 1900, *Termeszt. Füzetek*, XXIII, p. 317, Pl. VIII, figs. 7–8, ♀; 1911, *Ponerinae*, in ‘*Genera Insect.*,’ p. 90, ♀.

“WORKER.—Black; mandibles, antennae, legs and anus reddish; more densely sculptured with pubigerous punctures, the spaces between which on the head, thorax and petiole are opaque throughout; gaster less opaque; erect hairs absent even on the terminal abdominal segments. Head longer than broad, with feebly arcuate sides; eyes very minute, close to the mandibular articulation; clypeus moderately convex, its border broadly arcuate; mandibles shining, sparsely punctate, masticatory border anteriorly with three larger teeth, scarcely crenulate posteriorly. Antennal scape when folded back not reaching the occipital border; funiculi thick, all the joints except the first and last transverse, namely, 2–6 very short, 7 larger and together with the four last joints forming a 5-jointed club. Thoracic dorsum nearly straight, promesonotal suture distinct, mesoëpinotal suture obsolete: declivous surface of epinotum shining, abrupt, concave, acutely marginate on the sides, the angle be-