

is grayish and not very long, abundant only on the pleuræ, finer and rather sparse on the gaster. Legs, excluding the coxæ, castaneous, the tibiæ scarcely paler than the femora and not yellow as in the typical *labella* and the var. *obliqua*.

A single worker from the northwestern end of Bellona Island, Solomons (VI.21.33).

***Polyrhachis (Chariomyrma) rere* Mann**

Six workers from Anuda Island in the Santa Cruz group (VII.18.33). Originally described from Rere, on Guadalcanar Island, Solomons.

***Polyrhachis (Chariomyrma) kaipi* Mann**

A single worker from Anuda Island (VII.15.33). Originally described from Malaita Island, Solomons.

***Polyrhachis (Chariomyrma) arcuata* Le Guillou
var. *acutinoda* Forel**

One worker from the Kau Kau Plantation, Guadalcanar Island, Solomons (V.23.33). This variety was originally described from the Bismarck Archipelago.

***Polyrhachis (Hedomyrma) annæ* Mann**

Solomon Islands: Northwestern end of Bellona Island (VI.23.33) ♂—Kau Kau Plantation, Guadalcanar Island (V.20.33) ♀—Kungava Bay, Rennell Island (VI.14.33; VI.16.33) ♂ ♀.

Santa Cruz Islands: Mohawk Bay, Matema Island (VII.9.33) ♂ ♀.

Female (undescribed): Length 6–6.5 mm.

Resembling the worker; pronotal teeth shorter than broad at their bases and less acute; epinotal spines shorter and subparallel, petiolar spines straight and less divergent. Mesonotum as long as broad, semicircular in front. The dense golden pubescence covering the body the same as in the worker but somewhat less silvery on the head and thorax, the pale erect hairs slightly more numerous on the thorax and gaster.

Graciosa Bay, Santa Cruz Island is the type locality of this beautiful species, which was taken by Mann also in several localities in the Solomons.