

encouragement during the study. The research has been supported in part by National Science Foundation grant no. BSR 81-19350.

#### REFERENCES CITED

- BARONI URBANI, C.  
1980. The first fossil species of the Australian ant genus *Leptomyrmex* in amber from the Dominican Republic. *Stuttgarter Beitr. Naturkunde, Ser. B*, 62: 1-10.
- BROWN, W. L.  
1973. A comparison of the Hylean and Congo-West African rain forest ant faunas. In B. J. Meggers, E. S. Ayensu, and W. D. Duckworth, eds., *Tropical forest ecosystems: a comparative review*. Smithsonian Institution Press, Washington, D. C. Pp. 161-185.
- FOREL, A.  
1893. Formicides de l'Antille St. Vincent, récoltés par Mons. H. H. Smith. *Trans. Roy. Ent. Soc. London*, pt. 4, pp. 333-418.
- KEMPF, W. W.  
1959. A revision of the Neotropical ant genus *Monacis* Roger. *Studia Entomol. (n.s.)*, 2: 225-270.
- MAYR, G.  
1862. Myrmecologische Studien. *Verh. Zool.-bot. Ges. Wien*, 12: 649-766.
- WHEELER, W. M.  
1942. Studies of Neotropical ant-plants and ants. *Bull. Mus. Comp. Zool., Harvard*, 90: 1-262, 57 pls.
- WILSON, E. O.  
1985a. Ants of the Dominican amber (Hymenoptera: Formicidae). 1. Two new myrmicine genera and an aberrant *Pheidole*. *Psyche*, 92(1):1-9.  
1985b. Ants of the Dominican amber (Hymenoptera: Formicidae). 2. The first fossil army ants. *Psyche*, 92(1):11-16.