

Genus *ACANTHOLEPIS*.

Mayr, Europ. Formic. 1861.

8. *ACANTHOLEPIS FRAUENFELDI*, Mayr (Form. Austr. 1851), ♀.

This species has been hitherto known only from the Mediterranean country, Aden, Turkestan, and Persia. The specimens from Calcutta would belong to the variety *bipartita*, Smith (Proc. Lin. Soc. Zool. 1861); but the body is of a dark brown-blackish colour. They are large (3 millims.) and opaque.

The mesothorax of this genus is much narrowed in the middle, like the waist of a slender lady.

II. Subfamily *Dolichoderidæ*.

Forel, Zeitschr. f. wiss. Zool. xxx, Suppl. 1878.

No representative of this subfamily was among the ants sent from Calcutta.

III. Subfamily *Poneridæ*.

Smith, Cat. Brit. Hym. 1851.

Genus *PONERA*.

Latreille, Gen. Crust. et Ins. 1806-1809.

9. *PONERA TESSERINODA*, Mayr (Verhandl. k. k. zool.-bot. Ges. Wien, 1878, nec Emery, Ann. Mus. Civ. Genov. IX, 1876-77), ♀.

10. *PONERA LUTEIPES*, Mayr (Myrmecol. Studien, 1862), ♀.

Genus *DIACAMMA*.

Mayr, Myrmec. Studien, 1862.

11. *DIACAMMA VAGANS*, Smith (Journ. Linn. Soc. Zool. V, 1860, Roger (Berl. ent. Zeitschr. 1860, p. 304), ♀.

Genus *LOBOPELTA*.

Mayr, Myrmec. Studien, 1862.

12. *LOBOPELTA CHINENSIS*, Mayr (Neue Formiciden, 1870), ♀.

The specimen from Calcutta is only 8.5 millims. in length. Mayr's specimens were larger (9.5—10 millims.).

13. *LOBOPELTA KITTELI*, Mayr (Neue Formiciden, 1870), ♀.

IV. Subfamily *Dorylidæ*.

Shuckard, Ann. of Nat. Hist. v, 1840.

Genus *TYPHLOPONE*.

Westwood, Introd. Class. Ins. 1840.

14. *TYPHLOPONE ORANIENSIS*, Lucas (Expl. de l'Algérie, 1840-42), ♀.

This species has been hitherto known only from Northern Africa and from Asiatic Turkey. The specimens from Calcutta are quite identical with specimens from Algeria and Palestina in my collection.