

Indian Ants of the Indian Museum, Calcutta. No. 2.—By PROFESSOR AUGUST FOREL, Zurich. Communicated by the NATURAL HISTORY SECRETARY.

[Received May 20th ;—Read June 3rd, 1886.]

I. Subfamily **Camponotidæ**.

Genus CAMPONOTUS.

1. *C. SYLVATICUS*, Oliv., subsp. *C. cognatus*, Sm., var. e. Also found in Sibsagar (Assam) and in Sind.

2. *C. MICANS*, Nyl., ♂ and ♀. Variety of colour: the body, the legs, and the antennæ are partially or totally reddish brown or pale ferruginous. Sibsagar.

3. *C. GIGAS*, Latr. (Hist. nat. Fourmis), ♂. Johore, Malay Peninsula.

4. *C. SEXGUTTATUS*, Fab. (Ent. Syst. II. 354.)

Subsp. *C. exiguoguttatus*, n. st. This subspecies resembles also *C. cingulatus*, Mayr, but *cingulatus* has long, quite erect hairs on the tibiæ and scapi.

♀. Length 5.3—8.3 millims. Narrow, like the *C. sexguttatus*, i. sp., and *C. cingulatus*. Clypeus slightly carinate, very slightly and shortly lobate anteriorly, hardly emarginate in the middle of its anterior