



1. *Formica fusca* r. *lemani* Bondr. ♂
2. *Formica fusca* L. ♂ : a) the field among ocelli, b) the front part of pronotum-looking from above, c) squama petioli-looking from side.
3. The right half of phallus of *F. fusca* r. *lemani* Bondr.
4. Ninth abdominal sternum of *F. fusca* r. *lemani* Bondr.
5. The right half of phallus of *F. fusca* L.
6. Ninth abdominal sternum of *F. fusca* L.
7. Digitus and cuspis by *F. fusca* r. *lemani* Bondr. (a), and *F. fusca* L. (b).
8. Digitus and cuspis by *F. fusca* r. *lemani* Bondr. — variabilitas.

#### Description of male.

Material: Krkonoše 1946; Šumava 1947, 1948; Sobotka 1947  
Igt. Samšiňák.

Caput a little narrower than thorax. The composite eyes large, they project from the vault of head. Clypeus with a keel. Mandibulæ grooved along. Palpi mandibulares are longer but thinner than in the *Formica fusca* L.

Scapus overtops the crown about half its length. Funiculus fibrous, longer than the scapus. Its sections are swollen in first two third parts of its lower part. The first section of the funiculus is a little shorter, the last a little longer than the others. The field among ocelli carries besides the long bristles other short