

Additional Records. INDONESIA: Sulawesi: SW slope of Mount Klabat, 400–600 m, rotten wood in rainforest, nos. B9-1, K-2, K-61, X-9, W1-7, and Y-22, 13–19/VI/1972, 24 minor workers and nine majors (W. L. Brown, Jr., MCZ and BMNH); Air Madidi slope of Mount Klabat, 400–600 m, rotten wood in wet forest, nos. H3-6 and L-8, 13–19/VI/1972, 13 minor workers and two majors (W. L. Brown, Jr., MCZ and BMNH); Tangkoko-Batuangus Reserve, 200 m, nesting in log in rainforest, 29 July 1983, 37 minor workers and three majors (M. W. Moffett and D. A. Fletcher, MCZ, BMNH, MCSN and MHN). Seram: above Haruru, near Masohi, 50–150 m, rotten wood in rainforest, no. M-122, 18 March 1981, 12 minor workers (W. L. Brown, Jr., MCZ and BMNH). Seram minor specimens tending to be somewhat smaller than those from Sulawesi (HW averaging $[\bar{X} \pm SE]$ 1.13 ± 0.04 mm, versus 1.02 ± 0.04 mm for the Sulawesi minors; samples of 36 and 7 minors, respectively).

Natural History. The ants nest in cavities within logs in primary rainforests. Nests typically contain stores of fig seeds and other tiny seeds (see Moffett, 1985).

ACKNOWLEDGMENTS

I am grateful to W. L. Brown, Jr. and E. O. Wilson for critically reading the manuscript; B. Bolton, C. Besuchet, and R. Poggi for arranging loans of specimens and E. Seling for technical assistance with the SEM. This research was supported with grants from the National Geographic So-

ciety, the National Academy of Sciences, and Harvard University.

LITERATURE CITED

- EMERY, C. 1893. Formicides de l'ile de Ceylan. Ann. Soc. Ent. Fr., **62**: 239–255.
- . 1909. Expedition de la Nouvelle-Guinea 1904–05. Nova Guinea, **8**: 249–259.
- . 1922. Hymenoptera. Family Formicidae, subfamily Myrmicinae. In P. Wytsman (ed.), Genera Insectorum, fasc. 174, V. Verteneuil and L. Desmet, Brussels. 397 pp., 7 pl.
- HARRIS, R. A. 1979. A glossary of surface sculpturing. Occasional Papers in Entomology, California Dept. Food and Agriculture, no. 28, 31 pp.
- HENDY, M. D., AND D. PENNY. 1982. Branch and bound algorithms to determine minimal evolutionary trees. Math. Biosc. **59**: 277–290.
- KUGLER, C. 1978. A comparative study of the myrmicine sting apparatus. Studia Ent., **20**: 413–548.
- MOFFETT, M. W. 1985. Behavioral notes on the Asiatic harvesting ants *Acanthomyrmex notabilis* and *A. ferox*. Psyche, **92**: 165–179.
- SMITH, F. 1860. Descriptions of new species of hymenopterous insects collected by Mr. A. R. Wallace at Celebes. J. Proc. Linn. Soc. London Zool., **5**: 57–136.
- WHEELER, G. C., AND J. WHEELER. 1954. The ant larvae of the myrmicine tribe Myrmecinini. Proc. Entomol. Soc. Washington **56**(3): 126–138.
- . 1977. Supplementary studies on ant larvae: Myrmicinae. Trans. Amer. Entomol. Soc., **103**: 581–602.
- . 1983. Supplementary studies on ant larvae: Myrmicinae. Trans. Amer. Entomol. Soc., **108**: 601–610.
- WHEELER, W. M. 1919. The ants of Borneo. Bull. Mus. Comp. Zool. Harvard, **63**(3): 43–147.
- . 1930. Formosan ants collected by Dr. R. Takahashi. Proc. New England Zool. Club, **11**: 93–106.
- WILSON, E. O. 1985. Ants of the Dominican amber (Hymenoptera: Formicidae). I. Two new myrmicine genera and an aberrant *Pheidole*. Psyche, **92**(1): 1–9.