

latae, sub-laeyes aut plus minusve striolatae punctis dispersissimis; caput rotundato-quadrangulare, paulo longius quam latius, subtilissime coriaceum et subtiliter, disperse punctulatum; clypeus haud carinatus, deplanatus, margine antico arcuato; area frontalis in Operaria majori subtiliter, in Operaria minore haud impressa; thorax dense punctato-coriaceus, supra suturis leviter impressis; metanoto transversim rotundato, haud compresso, parte basali in medio impressione levi transversa; petioli squama incrassata, haud magna, antice convexa, postice plana, margine incrassato; abdomen densissime et subtilissime transverse striatum.

Haec species appropinquat generi: *Colobopsis*, capitis rotundato-rectangularis forma, clypeo parum convexo nec non thoracis forma.

9. *C. singularis* SMITH.

Formica singularis SM. *Cat.* 1858, pag. 27.

Camponotus singularis ROC. *Verz.* 1863, pag. 3.

Camponotus senilis MAYR *Myrm. Stud.* 1862, pag. 27.

Item in Java, Sumatra et Singapore.

Polyrhachis SHUCKARD.

History of Insects 1840.

Hoplomyrmus GERST. *Mon. ber. preuss. Akad.* 1858, PETERS *Reise. n. Mosamb. Zool., Ins.* pag. 507.

Hemioptica ROC. *Berl. ent. Z.* 1862, pag. 238.

10. *P. rastellata* LATREILLE.

Formica rastellata LTR. *Hist. nat. Fourm.* 1802, pag. 430.

Polyrhachis rastellata MAYR *Myrm. Stud.* 1862, pag. 40.

Polyrhachis busiris SMITH *Journ. Proc. Linn. Soc. V. Suppl.* 1860, pag. 98.

Polyrhachis euryalus SM. *Proc. Linn. Soc.* 1863, pag. 46.

Etiam in Ceilon, Celebes, Bouru, Amboina, Ceram, Mysole, Waigiou et Bachian.