

This variety is distinct from the typical form in color and somewhat larger size. The cleft in the upper border of the node is deeper than in the typical form, but this character is variable and I have only a small series of the latter to compare.

***Myrmelachista ambigua* Forel subspecies *ramulorum* Wheeler**

WHEELER, 1908, Bull. Amer. Mus. Nat. Hist., XXIV, p. 155, ♀, ♀, ♂.

Porto Rico:—Maricao; Mayaguez.

St. Thomas:—Charlotte Amalie; (Morrison).

***Camponotus (Myrmobrachys) planatus* Roger**

ROGER, 1863, Berliner ent. Zeitschr., VII, p. 146, ♀, ♀, ♂.

Cuba:—Occurs very abundantly through the island, where I found it in all localities visited, except in the vicinity of Pinares in the pine forest.

***Camponotus (Myrmoturba) santosi* Forel**

FOREL, 1908, Verh. zool-bot. ges. Wien, LVIII, p. 408, ♀.

Cuba:—San Antonio de los Banos; San Blas; Baracoa; Pinares; Mina Carlota; Limones.

***Camponotus (Myrmoturba) ramulorum* Wheeler**

WHEELER, 1905, Bull. Amer. Mus. Nat. Hist., XXI, p. 114, ♀, ♀.

Cuba:—Cienfuegos; Simpatia; Yateras District; San Blas; Cardenas; Victoria de las Tunas; Mina Carlota.

***Camponotus (Myrmoturba) maculatus* (Fabr.) subspecies *lucayanus*  
Wheeler**

WHEELER, 1905, Bull. Amer. Mus. Nat. Hist., XXI, p. 112, 3 ♀♀, ♀.

Bahamas:—Andros (Fresh Creek); Eleuthera (Bluff).

***Camponotus (Myrmeurynota) gilviventris* Roger**

ROGER, 1863, Berliner ent. Zeitschr., VII, p. 145, ♀.

WORKER MAJOR.—Length 7–7.5 mm.

Head slightly longer than broad, a little narrowed in front, with feebly convex sides, rather prominent and very narrowly rounded occipital corners and nearly straight border. Mandibles thick, with five strong, triangular teeth. Clypeus broader than long, with anterior border very shallowly concave at middle; surface convex at middle and shallowly depressed at sides anterior to middle. Frontal carinae strong, extending half the distance to occipital border. Frontal area triangular. Cheeks with a strong pit touching marginal suture of clypeus a little in front of middle. Eyes large and rather flat, situated in front of sides of head at posterior third. Antennal scapes arcuate, thickened distally, surpassing occipital corners by a distance equa-