

Described from a single worker taken from the stomach of a toad (*Bufo valliceps* Weig., U. S. N. M. 46967) collected by Nelson and Goldman, July 28, 1894.

The type lacks the antennal funiculi and portions of the legs, but is otherwise in good condition, with the pilosity well preserved. *Alfaria simulans* Emery, the nearest species, differs in its shorter and stouter antennal scapes, in the entire absence of an impression between meso- and epinotum, in its larger size and lighter color.

Leptogenys (Leptogenys) peninsularis, new species.

Worker.—Length 7.5 mm.

Head, excluding mandibles, more than one and one half times as long as broad, a little broader in front than behind, with feebly arcuate sides, broadly rounded occipital corners and straight border. Clypeus carinate at middle, the anterior projecting triangular portion as broad as long and subacuminate at tip, the median lobes at middle of sides of front border low and rounded. Mandibles distinctly shorter than sides of head, seen from the front nearly straight, broadened apically, inner border of basal portion subcultrate and terminating in an inner small, broad lamellate triangle, blades strongly concave. Antennal scapes surpassing occipital angles by about three-eighths of their length, second funicular joint one and one-half times as long as the first, remaining joints shorter and slightly thicker toward apex, the terminal as long as the two preceding joints together. Eye about as long as its distance to border of clypeus. Pronotum as broad as long, broadest behind middle, sides convex, dorsal surface slightly convex. Mesonotum slightly longer than broad, with straight sides and posterior border. Meso-epinotal impression moderate. Epinotal base nearly straight in outline and about twice as long as the flat declivous portion, into which it broadly rounds. Petiolar node higher than long, anterior surface broadly convex, posterior surface flat, dorsum broadly rounding into anterior and narrowly into posterior surface; from above longer than broad, slightly widest behind, with sides feebly arcuate. Abdomen elongate; sting strong. Legs long and slender.