

length 0.91 mm; thorax length 1.49 mm; pronotum width 0.67 mm; hind femur length 0.99 mm; petiole length 0.40 mm; petiole width 0.43 mm; postpetiole length 0.64 mm; postpetiole width 0.77 mm. Dark ferruginous-red; head and appendages slightly lighter in color.

Head shown in Figs. 16, 13. Mandibles dorsally smooth and shining, laterally superficially punctate; chewing border with approximately 8 larger triangular teeth, between the basal teeth there are minute intercalary denticles. Clypeus oblique in side view; anterior border convex in the middle. Frontal lobes 0.24 mm wide at their maximum expansion. Frontal sulcus reaching back to beyond posterior orbit of eyes. Scape finely punctate yet quite shining, its tip surpassing in repose the occipital border by 1 1/2 times its maximum thickness. Funicular segments I, VII-XI longer than broad, II-VI scarcely longer than broad to somewhat transverse. Sides of head gently convex, more constricted in front than behind; occipital border not excised. Dorsum of head subopaque, sharply and densely punctured; sculpture becoming superficial on sides, fading out completely on disc of gular face which is highly shining and smooth. Eyes relatively very large with about 8 facets across the greatest diameter.

Thorax as shown in Fig. 16. Promesonotum conspicuously convex in both directions. Antero-inferior corners of pronotum broadly rounded. Promesonotal suture impressed. Mesonotum almost as broad as long, distinctly bulging, with well delimited and impressed anterior, posterior and lateral borders. Mesoepinotal constriction pronounced both in profile and as seen from above. Basal face of epinotum transversely convex, not forming a narrow longitudinal ridge; lateral borders of declivous face bluntly marginate. Epinotal spiracle oval, facing obliquely caudad. Entire thorax, with the exception of the highly polished and smooth declivous face, densely rugulose-punctate and subopaque; mesopleura shinier due to vanishing sculpture on disc. Bottom of posterior portion of sides of thorax with fine horizontal striulae. Legs finely and superficially punctate, quite shining.

Petiole as shown in Fig. 16, rather stout, reticulate-rugose and punctate, sculpture vanishing on somewhat shining anterior and posterior face. Anterior face vertical below, oblique above, posterior face flat. Antero-basal tooth of node prominent. Subpetiolar process rounded. Gaster strongly constricted between first and second segment. Tergites I and II dorsally strongly,