

The writer is thankful to the Conselho Nacional de Pesquisas for financing this study under a fellowship, and to many collectors and friends who have enriched his collection (WWK), notably Fr. Thomas Borgmeier, O.F.M., Prof. W. L. Brown, Jr. of Cornell University; Mr. Carlos Alberto Campos Seabra, Rio de Janeiro; Prof. C. R. Gonçalves, of the Universidade Rural, Rio de Janeiro; Mr. K. Lenko, of the Instituto Biológico de São Paulo; Prof. Vicente Pérez D'Angello, of the Museo Nacional de Historia Natural de Santiago, Chile; Mr. Fritz Plaumann, of Nova Teutônia, Santa Catarina, Brasil; Prof. W. Weyrauch, of Lima, Peru.

In citing measurements and proportions, abbreviations standard in Brown's and Kempf's work are used:

Head length (HL), maximum measurable length of head proper as seen from dorsal full-face view, including all of clypeus and occipital lobes.

Head width (HW) is the maximum width of the cranium measured in the same view as for HL (includes also the eyes, when protruding laterally).

Scape length (SL) is the chord length of the antennal scape.

Weber's length (WL), oblique length of thorax (alitrunk) from side view, measuring from base of anterior pronotal declivity to metasternal extremity.

Hind femur length (HfL) is the chord length of the hind femur.

Total length (TL) of the body is the summed length of HL (plus closed mandibles), WL, plus the axial lengths of petiole, postpetiole and gaster measured separately.

Cephalic index (CI), head width expressed as a percentage of head length, or $HW/HL \times 100$.