

Santschi (1907) observed in an artificial nest, a wingless male of *C. nuda* var. *mauritanica* carrying larvae in a way similar to the worker's activity. This behavior was never observed by Lupo and Galil in nests of *C. wroughtoni* despite more than a year of observations (personal communication).

ACKNOWLEDGEMENTS

The author wishes to thank the following scientists and institutions for allowing him to study material under their care.

Dr. Q. Argaman, Dept. of Plant Protection, Tel Aviv; Dr. C. Baroni-Urbani (NHMB); Dr. C. Besuchet (MHNG); Mr. B. Bolton (BMNH); Dr. X. Espadaler, Universita Autonoma de Barcelona, Bellaterra; Prof. J. Galil (TAU); Dr. I. Löbl (MHNG); Dr. A. Lupo (TAU); Dr. R. Poggi (MSNG); Dr. B. Poldi, Mantova; Dr. R. Snelling (LACM); Mme. J. Weulersse (MNHP); Prof. E. Wilson (MCZ). Thanks are due to Mrs. T. Feler for technical help and Mr. W. Ferguson for drawing the figures.

REFERENCES

- André, E. 1881. Catalogue raisonné des Formicides provenant du voyage en Orient de M. Abeille de Perin et description des espèces nouvelles. *Annales de la Société Entomologique de France*, 1881:53-78 + 17 figs.
- André, E. 1881-1882. Species des Formicides d'Europe. In: *Species des Hyménoptères d'Europe et d'Algérie*, T. 2, 438 pp + 25 pls. Gray (Haute-Saône).
- Arnold, G. 1916. A monograph of the Formicidae of South Africa, part 2. *Annals of the South African Museum*, 14:159-270.
- Baroni-Urbani, C. 1971. Catalogo delle specie di Formicidae d'Italia. *Memorie della Società Entomologica Italiana*, 50:5-287.
- Baroni-Urbani, C. 1973. Die Gattung *Xenometra*, ein objektives Synonym (Hymenoptera, Formicidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, 46(3-4):199-201.
- Bernard, F. 1956. Révision des Fourmis paléarctiques du genre *Cardiocondyla* Emery. *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord*, 47:299-306 + 6 figs.
- Bernard, F. 1957. *Xenometra* Emery, genre de fourmis parasite nouveau pour l'Ancien Monde (Hym. Formicidae). *Bulletin de la Société Entomologique de France*, 62:100-103.
- Bernard, F. 1968. Les fourmis (Hymenoptera, Formicidae) d'Europe occidentale et septentrionale. Faune de l'Europe et du Bassin Méditerranéen, 3: Paris, Masson Ed., 411 pp., 425 figs.
- Bolton, B. 1982. Afrotropical species of the myrmicine ant genera *Cardiocondyla*, *Leptothorax*, *Melissotharsus*, *Messor* and *Cataulacus* (Formicidae). *Bulletin of the British Museum (Natural History), Entomology Series* 45(4):307-370.
- Borgmeier, T. 1937. *Cardiocondyla emeryi* Forel no Brasil, e a descoberta do macho ergatoide desta espécie (Hym. Formicidae). *Revista de Entomologia* 7(2-3):129-134.
- Brown, Jr., W.L. 1957. Is the ant genus *Tetramorium* native in North America? *Breviora, Museum of Comparative Zoology* 72:1-8.
- Creighton, W.S. 1950. The ants of North America. *Bulletin of the Museum of Comparative Zoology at Harvard College* 104, 585 pp. + 57 pls.
- Creighton, W.S. and R.R. Snelling. 1974. Notes on the behavior of three species of *Cardiocondyla* in the United States (Hymenoptera: Formicidae). *New York Entomological Society* 82:82-92.
- Donisthorpe, H. 1930. *Cardiocondyla bicolor* sp. n. (Hymenoptera, Formicidae), a species of Myrmecine ant new to science. *Annals and Magazine of Natural History*, Serie 10, 5:366.