

66. Anterior clypeal margin of the major sinuate; pronotum smooth and shining *bicarinata* subsp. *paiute*
 Anterior clypeal margin of the major bluntly bidentate; pronotum punctate and usually with transverse rugae or striae 67
67. Basal face of the epinotum in the major largely covered with transverse striae, and with punctures confined to the region of the meso-epinotal suture; pronotal rugae coarse and prominent *bicarinata*
 Basal face of the epinotum in the major largely punctate, transverse striae, when present, restricted to the area between the bases of the epinotal spines; pronotal rugae feeble 68
68. Epinotum of the minor armed with thick, short spines *bicarinata* subsp. *vinelandica*
 Epinotum of the minor armed with angular teeth which are broad at the base and do not resemble spines *bicarinata* subsp. *longula*

Subgenus CERATOPHEIDOLE

1. *Pheidole (Ceratopheidole) clydei* Gregg
 Ph. (Ceratopheidole) clydei Gregg, Jour. N. Y. Ent. Soc., 1950, 58, p. 89, ♀; Gregg, Amer. Mus. Novit., 1953, No. 1637, 4.
 Type locality: Carrizozo, New Mexico
2. *Pheidole (Ceratopheidole) grundmanni* M. R. Smith
 Ph. (Ceratopheidole) grundmanni M. R. Smith, Jour. N. Y. Ent. Soc., 1953, 61, p. 143, ♀.
 Type locality: Vernal, Utah

Subgenus MACROPHEIDOLE

3. *Pheidole (Macropheidole) rhea* Wheeler
 Ph. rhea Wheeler, Bull. Amer. Mus. Nat. Hist., 1908, 24, p. 452, ♀.
 Ph. (Macropheidole) rhea M. R. Smith, Proc. Ent. Soc. Wash., 1943, 45, p. 5, 4, ♀; Gregg, Psyche, 1949, 56, p. 70, ♀, ♀, 4; Creighton, Bull. Mus. Comp. Zool., 1950, 104, p. 168.
 Type locality: Nogales, Arizona

Subgenus PHEIDOLE

4. *Pheidole anastasii* Emery
 Ph. anastasii Emery, Bull. Soc. Ent. Ital., 1896, 28, p. 76, ♀, 4; Forel, Mitt. Naturh. Mus. Hamburg, 1901, 18, p. 78, ♀; Creighton, Bull. Mus. Comp. Zool., 1950, 104, p. 169.
 Type locality: Jimenez, Costa Rica