

The taxon *L. risi armouri* Wheeler, 1927 was described on a single worker and has never been found from then; the description matches very well a small worker of *L. risi* Forel but since we have not seen the type we prefer to wait for more material from its type locality, Barranco de la Virgen. Azuaje.

### References

- Barquín, J. 1981. Las hormigas de Canarias (Taxonomía, ecología y distribución de los Formicidae). Monogr. Univ. La Laguna, 3:1-586.
- Bhatkar, A.P. and W.H. Whitcomb, 1970. Artificial diet for rearing various species of ants. Flo. ent., 53:229-232.
- Bolton, B. 1987. A review of the *Solenopsis* genus-group and revision of Afrotropical *Monomorium* Mayr (Hymenoptera, Formicidae). Bull. Br. Mus. nat. Hist. (Ent). 54:263-452.
- Forel, A. 1892. Quelques fourmis de la faune méditerranéenne. Ann. Soc. Ent. Belg. 36:452-457.
- Santschi, F. 1909. *Leptothora rottenbergi* et especes voisines. Rev. Suisse Zool. 17:459-482.
- Wellenius, O. 1955. Formicidae Insularum Canariensium. Soc. Sci. Fennica. Comment. Biol. 15, 8:1-20.
- Wolf, H. 1980. Zur Kenntnis der Aculeaten-Fauna (Hymenoptera) von Gran Canaria und Teneriffa. Vieraea 9:65-78.

*Manuscript received on July 1987.*