

southeastern Brasil. All foraging *Stegomyrmex* workers found up to now have been seen collecting during this period, which we have called the favorable season.

Acknowledgements

We would like to thank Johan Billen for this expertise, the preparations, and photographs in Figures 1 a and 1 b. Anisio M. Diniz and Adelino D. da Silveira helped in the field. Bodo Dietz translated the German reference. Eliana M. Cancello, Flávia Ejchel, Márcia F. L. Françoso, Miriam D. Marques, Ricardo V. S. Paiva, Adriana R. Menezes, and specially P. E. Vanzolini critically read the manuscript. Claudio M. Martins identified the gastropods. CNPq, CAPES and FAPESP supported our work.

References

- Bolton, B., W. H. Gotwald and J.-M. Leroux, 1976. A new West African ant of the genus *Plectroctena* with ecological notes (Hymenoptera: Formicidae). *Ann. Univ. Abidjan sér. E (Ecologie)* 9:371–381.
- Brandão, C. R. F., J. L. M. Diniz and E. M. Tomotake, 1991. *Thaumatomyrmex* strips millipedes for prey: a novel predatory behaviour in ants, and the first case of sympatry in the genus (Hymenoptera: Formicidae). *Ins. Soc.* 38:335–344.
- Brown Jr., W. L., 1957. Predation of arthropod eggs by the ant genera *Proceratium* and *Discothyrea*. *Psyche* 64(3):115.
- Brown Jr., W. L., 1958. Contributions toward a reclassification of the Formicidae. II. Tribe Ectatomminae (Hymenoptera). *Bull. Mus. Comp. Zool. Harvard* 118(5):175–362.
- Brown Jr., W. L., 1974. A remarkable new island isolate in the ant genus *Proceratium* (Hymenoptera: Formicidae). *Psyche* 81(1):70–83.
- Brown Jr., W. L., 1979. A remarkable new species of *Proceratium*, with dietary and other notes on the genus (Hymenoptera: Formicidae). *Psyche* 86(4):337–346.
- Dejean, A. and J. P. Suzzoni, 1990. Monomorphism and polymorphism in African Ponerinae with a specialized alimentary diet: does this have any bearing on the type of prey? In: Veeresh, G. K., B. Mallik and C. A. Viraktamath (eds.) *Social Insects and the Environment*. Proc. 11th Int Congr. IUSSI, Bangalore. New Delhi, Oxford & IBH Publ. Co. Ltd, p. 339.
- Diniz, J. L. M., 1990. Revisão sistemática da tribo Stegomyrmicinae, com a descrição de uma nova espécie (Hymenoptera, Formicidae). *Revta bras. Ent.* 34(2):277–295.
- Eidmann, H., 1936. Oekologisch-faunistische Studien an südbrasilianischen Ameisen. *Arbeit. phys. angew. Ent. Berlin-Dahlem* 3(1):26–48, 81–114.
- Hölldobler, B. and E. O. Wilson, 1986. Soil-binding pilosity and camouflage in ants of the tribes Basicerotini and Stegomyrmecini (Hymenoptera, Formicidae). *Zoomorphology* 106:12–20.
- Hölldobler, B. and E. O. Wilson, 1990. *The Ants*. Cambridge, Belknap Press of the Harvard Univ. Press, 732 p.
- Lenko, K., 1965. Sôbre a ocorrência de *Stegomyrmex manni* no Estado de S. Paulo, Brasil (Hymenoptera: Formicidae). *Studia ent.* 8:201–204.
- Lévieux, J., 1972. Le rôle des fourmis dans les réseaux trophiques d'une savane préforestière de Côte-d'Ivoire. *Ann. Univ. Abidjan ser. E (Ecologie)* 5(1):143–240.
- Lévieux, J., 1982. A comparison of the ground dwelling ant populations between a Guinea savanna and an evergreen rain forest of the Ivory Coast. In: Breed, M. D., Michener, C. D. & Evans, H. E. (eds.) *The Biology of Social Insects*. Proc. 9th Int Congr. IUSSI, Boulder, Colorado, Westview Press, p. 48–53.
- Lévieux, J., 1983. The soil fauna of tropical savannas, IV: The Ants. In: Bourlière, F., (ed.) *Ecosystems of the World. 13. Tropical Savannas*, Amsterdam, Elsevier Scient. Publ. Co., p. 525–540.
- Villet, M. H., 1991. Colony founding in *Plectroctena mandibularis* F. Smith, and the evolution of ergatoid queens in *Plectroctena* (Hymenoptera: Formicidae). *J. Nat. Hist.* 25:979–983.