

16. *Acanthomyops* (*Chthonolasius*) *hingstoni*, sp. n.
(See below.)

17. *Camponotus paria*, Emery.

1 ♀, 2 ♂ ♂, Sikkim, Kalimpong, 4000 ft., 27. iii. 24.

18. *Camponotus badius*, Smith.

30 ♀ ♀ and ♂ ♂, N. India, Darjeeling, 5000 ft., 11-20. iii. 24.

19. *Camponotus nicobarensis*, Mayr.

1 ♀, Sikkim, Kalimpong, 4000 ft., 26. iii. 24.

20. *Camponotus wasmanni*, Emery.

1 ♂, Sikkim, Kalimpong, 4000 ft., 27. iv. 24.

21. *Polyrhachis dives*, Smith.

1 ♂, Sikkim, Kalimpong, 4000 ft., 27. iii. 24.

22. *Polyrhachis mayri*, Roze.

1 ♂, Sikkim, Kalimpong, 4000 ft., 26. iii. 24; 4 ♂ ♂, Sikkim, Gantok, 5000 ft., 3. iv. 24; 4 ♂ ♂, Sikkim, Singhik 4000 ft., 23. iv. 24.

Acanthomyops (*Chthonolasius*) *hingstoni*, sp. n.

♂. Reddish yellow, mandibles redder, teeth of mandibles and eyes black; whole body finely shagreened but shining, furnished with scattered, long, yellow hairs, and covered with fine, short, decumbent, yellow pubescence.

Head square, sides straight, posterior border straight when seen from front, slightly excavate when seen from behind; *mandibles* massive, finely striate, masticatory margin furnished with eight blunt teeth; *clypeus* ample, convex, subcarinate, slightly longer than broad, anterior border somewhat rounded in centre, and slightly crenulate; *frontal area* conspicuous though indistinctly defined; *ocelli* distinct; *frontal furrow* well marked, reaching median ocellus; *eyes* normal; *frontal carinæ* well marked; *antennæ* long, *scape* slightly curved at base, when bent back reaching beyond margin of head by nearly half its length, *funiculus* gradually thickened to apex, with all the joints longer than broad; *maxillary palpi* only 5-jointed (joints 5 and 6 evidently fused together); *labial palpi* 4-jointed.