

*eyes* large, placed laterally and rather forward on head; *antennæ* long, *scape* slightly curved, a little broadest before apex, when bent back reaching beyond posterior border of head by about one-third of its length; *funiculus* with joints elongate, 1st joint longer and a little broader than 2nd, 2nd-5th subequal, 6th-9th gradually decreasing in length and increasing in breadth, 10th a little longer than broad, last joint about once and a half as long as the one preceding. *Thorax* narrow; *pronotum* convex, rounded at sides; *epinotum* convex, globose, higher when seen in profile than mesonotum; *node* of *pedicel* long, pointed, inclined forward, flat anteriorly and posteriorly; *gaster* small, narrow-oval, not quite twice as long as broad. *Legs* long. *Long.* 3.2 mm.

Described from a single worker taken by Dr. Edwards at the eastern end of Lake Nahuel Huapi, 2000 ft., 28-31. x. 26. Type in Coll. Brit. Mus.

FORMICINÆ.

*Jaisiphane*

*Acanthomyops (Donisthorpea) edwardsi*, sp. n.

♂.—Yellowish brown, posterior part of head and gaster darker, eyes black; whole body smooth and shining (the front of head, mesonotum, and epinotum, however, very finely punctured when seen under a strong power); pubescence very sparse and minute, a few scattered hairs on head and gaster. *Head*, without mandibles, slightly longer than broad, somewhat rounded behind, with posterior border straight; *mandibles* long, striate, masticatory margin furnished with five rather sharp teeth, the last long and pointed, 2nd to 4th longer than 1st; *clypeus* broader than long, convex, distinctly carinate, anterior border straight (the carina from one point of view, however, gives it the appearance of being pointed); *frontal area* conspicuous, but with basal portion indistinctly defined; *ocelli* minute, but distinct; *frontal furrow* very faint; *eyes* oval, slightly prominent; *frontal carinæ* short but well marked; *antennæ* fairly long, *scape* without outstanding hairs, very slightly curved, when bent back reaching beyond margin of head by about one-fourth of its length; *funiculus*, 1st joint longer and broader than 2nd, 2nd-12th gradually increasing in breadth, 2nd slightly longer than 3rd, 3rd and 4th of about equal length, 5th-12th gradually increasing