

(2 winged ♀♀ were taken at L. Nahuel Huapi, Puerto Blest, 2500 ft., 2-3. xii. 26). Type and allotypes in Coll. Brit. Mus.

*Acanthomyops* (*Chthonolasius*) *negrensis*, sp. n.

♀.—Dark brown, shining, mandibles, clypeus, cheeks, antennæ and legs brownish yellow; clothed with fine decumbent yellowish-grey pubescence, and scattered yellow outstanding hairs; puncturation very fine and close. *Head* as broad at base as thorax; *mandibles* long, longitudinally striate, and with a few widely separated punctures, masticatory margin furnished with eleven pointed teeth, terminal tooth rather long, penultimate tooth longer than those before it, 1st tooth shortest, 2nd bifid; *clypeus* broad, convex, subcarinate; *frontal area* large and indistinctly defined; *frontal furrow* faint; *eyes* large, prominent; *front* with a small circular pit before median *ocellus*; *antennæ* long, *scape* with outstanding hairs, when bent back reaching beyond margin of head by about one-third of its length, 1st joint of *funiculus* longer and broader than 2nd, 2nd very slightly longer than 3rd, 3rd-11th (=12th joint of antenna) increasing in length and breadth, 11th about equal to the two preceding it taken together. *Thorax* broad, convex; *scale* broadest in middle, apex distinctly but not deeply emarginate; *gaster* broader than thorax, but not very voluminous. *Wings* slightly infuscate; *veins* and *pterostigma* yellow. *Tibiæ* with outstanding hairs. *Long.* 8.6 mm.

Described from a single winged ♀ taken by Dr. Edwards at L. Nahuel Huapi, Eastern End, 17. xi. 26. Type in Coll. Brit. Mus.

*Acanthomyops* (*Chthonolasius*) *rufo-niger*, sp. n.

♂.—*Head*, *thorax*, *scale*, and *scapes* of antennæ light brownish red, *eyes* black, *gaster* black, reddish at extreme base; *mandibles*, *legs*, and *funiculi* of antennæ brown, shining; *pubescence* fine and sparse; a few scattered yellow hairs, which are longer on *gaster*.

*Head* rather large, broadest before base, sides rounded, posterior border slightly rounded; *mandibles* massive, longitudinally striate, masticatory border armed with six blackish pointed teeth, the terminal one being the longest, the 3rd and 5th longer than the rest; *clypeus*