

*Distribution*.—India, Malaya, Hawaiian Isles, Borneo, New Guinea.

Some 60 specimens, including ♂♂, 1 deäl. ♀, 4 4, and ♀♀.

10. *Tetramorium guineensis* F.

*Formica guineense* Fabricius, Ent. Syst. ii. p. 357 (1793).

*Tetramorium guineensis* Donisthorpe, Bull. R. Bot. Gard. Kew, xii. p. 368 (1911); Emery, Gen. Ins. clxxiv. p. 278 (1922). For synonymy, see Donisthorpe, Brit. Ants, 2nd ed. p. 391 (1927).

*Distribution*.—Cosmopolitan species.

Two ♀♀.

11. *Tetramorium simillimum* Smith.

*Myrmica simillima* Smith, List Brit. Anim. Brit. Mus., Acul. p. 118 (1851).

*Tetramorium simillimum* Donisthorpe, Bull. R. Bot. Gard. Kew, iii. p. 122 (1908); Emery, Gen. Ins. clxxiv. p. 279 (1922); Donisthorpe, Brit. Ants, 2nd ed. p. 392 (1927).

*Distribution*.—Cosmopolitan species.

Two winged ♀♀, two deäl. ♀♀, 22 ♀♀.

12. *Triglyphothrix striatidens* Emery.

*Tetramorium obesum* razza *striatidens* Emery, Ann. Mus. Civ. Gen. xxvii. p. 501 (1889).

*Triglyphothrix striatidens* Donisthorpe, Bull. R. Bot. Gard. Kew, iii. p. 122 (1908); Emery, Gen. Ins. clxxiv. p. 274 (1922); Donisthorpe, Brit. Ants, 2nd ed. p. 393 (1927).

*Distribution*.—India, Burma, Ceylon, Tunis, Sierra Leone, etc., becoming cosmopolitan.

Two ♀♀.

*DOLICHODERINÆ.*

13. *Tapinoma melanocephalum* F.

*Formica melanocephalum* Fabricius, Ent. Syst. ii. p. 353 (1793).

*Tapinoma melanocephalum* Emery, Gen. Ins. cxxxvii. p. 41 (1912); Donisthorpe, Brit. Ants, 2nd ed. p. 395 (1927).

*Distribution*.—Cosmopolitan species.

21 ♀♀.

*FORMICINÆ.*

14. *Anoplolepis (Anoplolepis) longipes* Jerd.

*Formica longipes* Jerdon, Madras Journ. Litt. Sci. xxvii. p. 112 (1851).

*Plagirolepis longipes* Wheeler, Bull. Mus. Compar. Zool. lxiii. p. 366 (1919).

*Anoplolepis (Anoplolepis) longipes* Emery, Gen. Ins. clxxxiii. p. 17 (1928).