

Distribution.—Tropical Asia. Imported by commerce into tropics, Borneo, Solomon Isls., Fiji, etc. Becoming a cosmopolitan species: see Donisthorpe, Brit. Ants, 2nd ed. p. 385 (1927).

Sixteen ♂♂.

√ 15. *Camponotus (Tanæmyrmex) melichloros* Kirby.

Camponotus melichloros Kirby, Proc. Zool. Soc. Lond. p. 551 (1888);
Emery, Gen. Ins. clxxxiii. p. 172 (1925).

Distribution.—Christmas Island.

Over 150 specimens, including ♂♂, one deäl. ♀, 4 ♀, and
♂♂.

16. *Camponotus (Tanæmyrmex) sp. ?*

Two ♂♂.

17. *Paratrechina (Paratrechina) longicornis* Latr.

Formica longicornis Latreille, Hist. Nat. Fourm. p. 113 (1802).
Tapinoma gracilescens Smith, Cat. Hym. Brit. Mus. vi. p. 56 (1858).
Prenolepis longicornis Donisthorpe, Bull. R. Bot. Gard. Kew, iii.
p. 122 (1908).

Paratrechina (Paratrechina) longicornis Emery, Gen. Ins. clxxxiii.
p. 217 (1925); Donisthorpe, Brit. Ants, 2nd ed. p. 399 (1927).

Distribution.—Cosmopolitan species.

One winged ♀, one deäl. ♀, 143 ♂♂.

18. *Paratrechina (Nylanderia) vividula* Nyl.

Formica vividula Nylander, Acta Soc. Sc. Fenn. ii. p. 900 (1846).
Prenolepis vividula Donisthorpe, Bull. R. Bot. Gard. Kew, iii. p. 122
(1908).

Paratrechina (Nylanderia) vividula Emery, Gen. Ins. clxxxiii. p. 233
(1925); Donisthorpe, Brit. Ants, 2nd ed. p. 398 (1927).

Distribution.—Cosmopolitan species.

Two ♂♂, 8 ♂♂.

19. *Paratrechina (Nylanderia) bourbonica* For.

Prenolepis nodifera bourbonica Forel, Ann. Soc. Ent. Belg. xxx.
p. 210 (1866).

Paratrechina (Nylanderia) bourbonica Emery, Gen. Ins. clxxxiii. p. 219
(1925).

Distribution.—Reunion, Pembri Island, Madagascar, New Caledonia, Chigos Island, Solomon Atoll, Seychelles. Probably becoming cosmopolitan. Several subspecies have been described from Bengal, Hawayii Isles, Easter Island, etc.

5 ♂♂, 1 deäl. ♀, 32 ♂♂.