

broad, distinct, and longitudinally striate; rest of mesothorax behind mayrian furrows, and between parapasidal furrows, with scattered large shallow oblong punctures and some impressed lines, a deep longitudinal pointed furrow occurs outside parapasidal furrow; *episternite* and *epimera* of mesothorax and *episternite* of metathorax transversely striate, *furrow* between episternite and epimera of mesothorax deep, and widely longitudinally striate; *suture* between præscutellum and scutellum not defined; *præscutellum* consisting of a rather deep transverse hollow in centre, which is furnished with a number of widely separated raised longitudinal ridges; *scutellum* convex, prominent, smooth in centre, with shallow oblong punctures on anterior portion, and with rather deep hollows which are longitudinally striate at sides; *metanotum* transverse, narrow, smooth, convex, and prominent in centre, somewhat hollowed out and finely longitudinally striate at sides; *epinotum* longer than broad, narrowed to base, *dorsal surface* covered with a network of strongly-raised ridges, *declivity* flat hexagonal in shape, bordered by raised ridges, with a longitudinal raised ridge in centre from which raised transverse ridges branch. *Petiole* longer than broad with a high node rounded above, narrowed anteriorly with a small tubercle on each side before base and a very small blunt tooth beneath, upper surface smooth, sides covered with a network of closely-raised ridges, posteriorly somewhat triangularly excavated, the upper portion smooth, the lower with longitudinally-raised ridges; *post-petiole* smooth, pyriform in shape, considerably longer and broader than petiole, constriction between post-petiole and second segment of gaster very marked; *gaster* smooth, long oval, broadest before base of second segment; *pygidium* terminating in a fairly long curved sharply-pointed spine; *subgenital lamina* fairly long, bluntly pointed at apex; *cerci* present. *Wings* iridescent, *veins* and *pterostigma* yellow, one *discoidal cell* and two long *cubital cells* present, *radial cell* long, closed, appendiculate at apex. *Legs* long, slender, two *spurs* to posterior pairs of *tibiæ*, *claws* bidentate.

*Long.* 10.5 mm.

Described from a male taken by C. F. Baker at Mt. Banohao, Philippines, 1924—486.

Type in Brit. Mus. Coll.

*Ann. & Mag. N. Hist. Ser. 11. Vol. ix.*